



Therapeutic options in recurrent glioblastoma-An update

Seystahl, Katharina ; Wick, Wolfgang ; Weller, Michael

Abstract: INTRODUCTION Standards of care are not yet defined in recurrent glioblastoma. METHODS We reviewed the literature on clinical trials for recurrent glioblastoma available in PubMed and American Society of Clinical Oncology (ASCO) abstracts until June 2015. RESULTS Evidence is limited due to the paucity of randomized controlled studies. Second surgery or re-irradiation are options for selected patients. Alkylating chemotherapy such as nitrosoureas or temozolomide and the vascular endothelial growth factor (VEGF) antibody, bevacizumab, exhibit comparable single agent activity. Phase III data exploring the benefit of combining bevacizumab and lomustine are emerging. Novel approaches in the fields of targeted therapy, immunotherapy, and tumor metabolism are coming forward. Several biomarkers are being explored, but, except for O(6)-methylguanine DNA methyltransferase (MGMT) promoter methylation, none has assumed a role in clinical practice. CONCLUSION Proper patient selection, development of predictive biomarkers and randomized controlled studies are required to develop evidence-based concepts for recurrent glioblastoma.

DOI: <https://doi.org/10.1016/j.critrevonc.2016.01.018>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-124230>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Seystahl, Katharina; Wick, Wolfgang; Weller, Michael (2016). Therapeutic options in recurrent glioblastoma-An update. *Critical Reviews in Oncology/Hematology*, 99:389-408.

DOI: <https://doi.org/10.1016/j.critrevonc.2016.01.018>

Therapeutic options in recurrent glioblastoma – an update

Katharina Seystahl^{1*}, Wolfgang Wick², Michael Weller¹

¹Department of Neurology, University Hospital Zurich, Zurich, Switzerland

²Department of Neurology and Neurooncology Program National Center of Tumor Diseases, University Hospital Heidelberg, Heidelberg, Germany

Running title: Therapeutic options in recurrent glioblastoma

***Corresponding Author:**

Katharina Seystahl, MD

Department of Neurology

University Hospital Zurich

Frauenklinikstrasse 26, CH-8091 Zurich, Switzerland

Tel: (41) 44 2555500

Fax: (41) 44 2554507

Email: katharina.seystahl@usz.ch

Abstract

Introduction

Standards of care are not yet defined in recurrent glioblastoma.

Methods

We reviewed the literature on clinical trials for recurrent glioblastoma available in PubMed and American Society of Clinical Oncology (ASCO) abstracts until June 2015.

Results

Evidence is limited due to the paucity of randomized controlled studies. Second surgery or re-irradiation are options for selected patients. Alkylating chemotherapy such as nitrosoureas or temozolomide and the vascular endothelial growth factor (VEGF) antibody, bevacizumab, exhibit comparable single agent activity. Phase III data exploring the benefit of combining bevacizumab and lomustine are emerging. Novel approaches in the fields of targeted therapy, immunotherapy, and tumor metabolism are coming forward. Several biomarkers are being explored, but, except for *O*(6)-methylguanine DNA methyltransferase (MGMT) promoter methylation, none has assumed a role in clinical practice.

Conclusion

Proper patient selection, development of predictive biomarkers and randomized controlled studies are required to develop evidence-based concepts for recurrent glioblastoma.

Keywords: Glioblastoma, MGMT, temozolomide, nitrosourea, bevacizumab, immunotherapy

Highlights:

1. Standards of care are incompletely defined in recurrent glioblastoma.
2. The evidence for repeat surgery or reirradiation is limited.
3. Alkylating chemotherapy (nitrosourea, temozolomide) is a widely accepted therapeutic option.
4. Bevacizumab has clinical activity, but an effect on overall survival is uncertain.
5. Individualized treatment concepts should consider age, performance status, *MGMT* promoter methylation status, response to and type of previous regimens and quality of life.

Contents

1. Introduction
2. Scope and objectives
3. Results
 - 3.1 Diagnosis of progression and response
 - 3.2 Surgery at recurrence
 - 3.3 Repeat radiotherapy
 - 3.4 Chemotherapy for recurrent glioblastoma
 - 3.4.1 Nitrosourea monotherapy and combination regimens
 - 3.4.2 Temozolomide monotherapy and combination regimens
 - 3.4.3 Other classical non-alkylating chemotherapy
 - 3.4.4 Bevacizumab monotherapy and combination regimens
 - 3.4.5 Targeted therapy
 - 3.4.6 Immunotherapeutic approaches
 - 3.4.7 Alternative approaches
4. Summary and conclusion
5. Conflicts of interest statement
6. Tables
7. Vitae
8. References

1. Introduction

Despite advances in the understanding of its biology, glioblastoma, the most common malignant brain tumor, remains a devastating disease. Median overall survival (OS) in population-based studies in the US was 8.1 months in the period of 2000 – 2003 and 9.7 months from 2005 -2008 [1]. In 2005, the current standard of care in newly diagnosed glioblastoma was established based on the trial of the European Organisation for Research and Treatment of Cancer (EORTC)/National Cancer Institute of Canada Clinical Trials Group (NCIC-CTG) showing prolonged median OS of 14.6 months by addition of temozolomide (TMZ) during and after radiotherapy compared to radiotherapy alone (12.1 months) [2]. Promoter methylation of the O⁶-methylguanine-DNA methyltransferase (*MGMT*) gene was established as a predictive biomarker for benefit from TMZ in newly diagnosed glioblastoma [3]. For elderly patients, e.g., older than 65-70 years, the standard of care for newly diagnosed glioblastoma without *MGMT* promoter methylation or unknown *MGMT* status is radiotherapy alone [4]. In contrast, elderly patients with glioblastoma with *MGMT* promoter methylation should receive TMZ without or with radiotherapy [5-7]. Tumor recurrence occurs in almost all patients. Currently, no standard of care is established in recurrent or progressive glioblastoma [7]. Despite numerous clinical trials, the identification of effective therapies is complicated by the lack of appropriate control arms, selection bias, small sample sizes and disease heterogeneity [8].

2. Scope and objectives

Here we review clinical efficacy data for the treatment of patients with recurrent (=progressive) glioblastoma with focus on prospective and randomized clinical trials published in PubMed or as ASCO abstract reports until June 2015. Phase I or retrospective studies were only included in case of lack of prospective data, or for historical or innovative importance. We discuss the current status of diagnosis of tumor response and progression. The therapeutic value of repeat surgery and radiotherapy as well of different systemic treatment regimens including nitrosoureas, temozolomide, bevacizumab, targeted therapies, combinational approaches or novel concepts including immunotherapy in patients with recurrent glioblastoma are reviewed, with a focus on new data emerging since 2013 [8]. Finally, the impact of these clinical trials on current clinical practice in recurrent glioblastoma is discussed.

3. Results

3.1 Diagnosis of progression and response

Criteria for response and progression in glioblastoma have remained a matter of debate for the past decades. Still, MRI imaging every 2 to 3 months remains the gold standard for diagnosis of response or progression in glioblastoma. However, several concerns have been raised regarding the Macdonald criteria, being the standard tool for response evaluation until 2010 [9]. Limitations included the observation of “pseudoprogression” as a transient increase of contrast-enhancing tumor especially within the first 3 months after completion of radiotherapy as well as “pseudoresponses” with divergent effects on contrast-enhanced T1-images versus T2/fluid-attenuated inversion recovery (FLAIR) sequences especially with use of antiangiogenic agents. Since 2010, the updated criteria for Response Assessment in Neuro-Oncology (RANO) add restricted parameters for diagnosis of progressive disease within 3 months after completion of radiochemotherapy and integrate the evaluation of T2/FLAIR sequences as well of corticosteroid use [10]. Recently, the same group has provided an international consensus protocol based on the development of the EORTC Brain Tumor Group for uniform evaluation in clinical trials and practice [11]. Comparing overall response rates and progression-free survival (PFS) in different clinical trials using either Macdonald or RANO criteria is thus difficult. To date, the RANO criteria are considered the best standardized tool for evaluation of response and progression in glioblastoma. Several imaging techniques and modalities have been proposed for the identification of response or progression as well as prognostic or predictive tools for therapeutic efficacy, e.g. perfusion imaging, dynamic susceptibility contrast, apparent diffusion coefficient and MR spectroscopy [12-18]. However, these studies are limited in patient size and/or adequate control groups. To date, in case of suspected pseudoprogression, repeated MRI imaging in shortened time intervals is recommended, while usually maintaining treatment. O-(2-(18)F-fluoroethyl)-L-tyrosine)-positron emission tomography ((18)F-FET-PET) may help to identify pseudoprogression versus early progression after radiochemotherapy, too [19].

3.2 Surgery at recurrence

The role of repeat surgery in patients with progressive or recurrent glioblastoma remains controversial. Some retrospective studies proposed a survival benefit after reoperation [20-23] while others did not [24]. Limitations of most of the studies

include high probability of selection bias and lack reasonable controls for comparison [25]. Factors proposed for benefit of surgery and decision-making represent general prognostic factors for glioblastoma, such as age, *MGMT* promotor methylation, performance status, tumor volume, extent of resection, and tumor localization. A scale including tumor involvement in non-eloquent areas, small tumor volume (< 50 cm³) and good performance status (KPS>80%) was proposed in 2010 and modified to a score comprising KPS and ependymal involvement for decision making regarding reoperation, based on retrospective data [26, 27].

Few prospective data sets on repeat surgery are available. A prospective registry study of 764 patients with glioblastoma analyzed over a 7-year period (2004-2010) with one third of patients reoperated at recurrence did not suggest a benefit of repeat surgery (HR 1.02), however, no detailed patient stratification for prognostic variables was available [28]. Similarly, in a meta-analysis of 8 prospective phase I and II trials comprising 300 patients with recurrent glioblastoma repeat surgery was not an independent predictor for PFS and OS [29]. A post-hoc analysis of the prospective DIRECTOR trial evaluating two different dosing schemes of TMZ in patients with recurrent glioblastoma did not show a difference in overall survival in patients with (n=71) versus without (n=34) surgery at recurrence. However, when stratifying for extent of resection (n=59 available for analysis), there was superior survival in patients with complete resection of gadolinium-enhancing tumor (n=39) [30]. Similarly, a prospective non-randomized registry for patients with repeat surgery identified a correlation between (high) extent of resection and survival [31].

Beyond an expected therapeutic efficacy of repeat surgery, the acquisition of tumor tissue at recurrence might be valuable in some situations, e.g. for diagnosis of recurrent disease (e.g. versus radiation necrosis), confirmation of initial histology and/or determination of molecular markers for biomarker-based decision making.

3.3 Repeat radiotherapy

Re-irradiation is a similarly controversial option for patients with recurrent glioblastoma. Evidence-based clinical decision making is impeded by the lack of prospective and randomized trial design and high probability of selection bias in single arm studies. Main concerns include safety with regard to radiation necrosis and neurocognitive impairment as well as limited efficacy at recurrence after initial irradiation. Without establishment in prospective or controlled studies, for fractionated

stereotactic radiotherapy total doses between 30-36 Gy in 2-3.5 Gy fractions with or without intensity modulation have been used [32, 33]. Radiosurgery is less frequently used in glioblastoma. As favorable variables for selecting patients likely to benefit from re-irradiation again general prognostic factors such as young age, high KPS or small tumor size have been proposed [32, 34-37]. Different reports exist regarding the preferred time interval to initial radiation for benefit from re-irradiation, favoring either an interval of less than 6 months [35] or more than 6 or even 12 months [37, 38].

Several combination partners have been proposed for re-irradiation with the hypothesis of synergistic effects. A randomized phase II study evaluating re-irradiation with and without the addition of a CD95 ligand-binding fusion protein, APG101, showed poor outcome in the re-irradiation-only arm (PFS-6 3.8%, median PFS 2.5 months) but significantly higher PFS-6 (20.7%) and median PFS (4.5 months) in the combination arm. However, median OS was equal in both groups with 11.5 months. Presence of CD95L as determined by promoter methylation may identify patients benefitting from APG101 [39]. Several studies evaluated the combination of bevacizumab with re-irradiation with the hypothesis that vascular normalization results in improved oxygenation of tumor tissue and thereby increases the effect of radiation. With heterogeneous cohorts of WHO grade III (n=6, n=14, n=12) and glioblastomas (n=8, n=43, n=42) treated with re-irradiation with and without bevacizumab a higher median PFS (5.7 vs. 3.7, 5.6 vs. 2.5, and 6 vs. 4 months) and higher median OS (8.3 vs. 14.3, 8.6 vs. 5.7, 11 vs 8.3 months) were reached in 2 of 3 retrospective studies in the bevacizumab cohort [40-42]. In the study of Minniti and colleagues the non-bevacizumab group received fotemustine. A prospective study evaluating stereotactic radiosurgery plus bevacizumab (n=8 patients with anaplastic glioma, n=7 with glioblastoma) reported a median PFS of 3.9 and median OS 14.4 months. Toxicity included one grade 3 and two grade 2 CNS toxicities [43].

3.4 Chemotherapy for recurrent glioblastoma

The landscape of chemotherapeutic trials in recurrent glioblastoma is influenced by the historical evolution of the standard of care for newly diagnosed and recurrent disease, heterogeneity of inclusion criteria, choice of endpoints and response criteria, pretreatment and patient characteristics. In addition, most studies are non-

comparative or fail to use a control arm lacking the experimental drug.

Some, especially older trials include both anaplastic gliomas and glioblastoma. Trials conducted before the establishment of standard TMZ-based radiochemotherapy in 2005 commonly included TMZ-naïve patients. Different endpoints and use of different response criteria further impede comparability of trials. Six-months PFS rate (PFS-6) and median OS from the time of treatment for recurrence are reasonable endpoints for clinical trials in recurrent glioblastoma [44], however, criteria of response and progression are still moving targets.

3.4.1 Nitrosourea monotherapy and combination regimens

Nitrosoureas (e.g., carmustine [BCNU], lomustine [CCNU], nimustine [ACNU], fotemustine) are DNA alkylating agents that cross the blood-brain barrier and have been extensively used in glioma treatment. They may induce considerable hematological toxicity with long-lasting bone marrow suppression, liver and renal toxicity, and, specifically carmustine, interstitial lung disease. After approval of TMZ in 1999 for recurrent glioblastoma based on efficacy data of 2 phase II trials and a favorable safety profile [45, 46], nitrosoureas were used less frequently. Their use rose again when TMZ became standard of care in newly diagnosed glioblastoma. Table 1 summarizes nitrosourea-based trials in recurrent glioblastoma. There are 5 single arm phase II trials with nitrosourea monotherapy (1 with carmustine, 4 with fotemustine at different dosing regimens) [47-51] and 6 randomized phase II or phase III trials comprising one therapeutic arm with nitrosourea monotherapy: 5 trials with lomustine [52-56] and 1 trial with carmustine where data for patients treated with carmustine (n=29) were not separately reported but summarized in a common “control” arm with 27 patients receiving TMZ [57]. All but 3 trials were conducted at the first recurrence after TMZ-based standard radiochemotherapy. One study was done in chemotherapy-naïve-patients [47], one study did not disclose previous chemotherapy except exclusion of nitrosourea, bevacizumab or investigational agents [53], one study reported previous chemotherapy in about 60% of the patients without details [57]. PFS-6 ranged between 17.5% and 61.5% and median OS between 6.0 and 11.1 months for monotherapy of nitrosourea agents. Notably, in the randomized studies, lomustine as monotherapy showed comparable results with the investigational agents enzastaurin, cediranib, galunisertib or bevacizumab [52-55], pointing towards relevant single agent activity of the “control”

agent or lack of efficacy of the experimental agents. The combination of lomustine plus bevacizumab showed prolonged median PFS and OS and higher PFS-6 than the single agents in the BELOB phase II trial [54]. However, the combination treatment at least with the higher dose of lomustine with 110 mg/m² exhibited more hematological toxicity leading to the dose reduction to 90 mg/m² when combined with bevacizumab. Another randomized phase II trial evaluated the combination of lomustine with 90 mg/m² with bevacizumab 5 mg/kg every 3 weeks versus bevacizumab alone at 10 mg/kg every 2 weeks in a heterogeneous cohort of patients with glioblastoma at 1st, 2nd or 3rd relapse. The authors emphasized a benefit for the subgroup of patients at 1st relapse in the combination arm with a median OS with 13.1 months versus 8.8 months in the monotherapy arm without disclosing the number of patients in this subgroup and without reporting PFS and OS results of the intent-to-treat population [58]. Another phase II study evaluated the combination of nitrosourea, namely fotemustine with bevacizumab, however, not in a randomized but single-arm design [59]. PFS-6-rate of 42.6% was similar with the BELOB trial with a median OS of 9.1 months (BELOB: 11 months [54]). Again, hematological toxicity was relevant with 22% of the patients discontinuing fotemustine due to grade 3 or 4 myelotoxicity. The promising efficacy signal of the combination of nitrosoureas with bevacizumab of phase II trials was not confirmed in the EORTC 26101 phase III trial comparing lomustine plus bevacizumab with lomustine alone in patients with recurrent glioblastoma which did not report a difference in OS (8.6 versus 9.1 months), although prolonged PFS (1.5 versus 4.2 months) was confirmed [56]. Other studies regarding nitrosourea combination therapies for recurrent disease are mostly retrospective. One phase II trial evaluating the combination of carmustine and irinotecan showed PFS-6 of 30.3%, median PFS of 3.9 and median OS of 11.7 months. Beyond hematologic toxicities, mainly grade 1 and 2, frequent diarrhea (69% grade 1 and 2, 7.1% grade 3) and 9.5% cholinergic syndrome were seen [60]. The combination of TMZ with nitrosoureas, fotemustine or carmustine, was evaluated in 2 prospective studies, both with severe and frequent hematologic toxicities leading to dose reductions and early stop of enrolment in one study without clear efficacy signals beyond single agent activity [61, 62].

In summary, the different nitrosoureas show probably comparable efficacy as single agents and remain to be one standard of care at least for current clinical trials. It is likely to expect that clinical efficacy will be more prominent in patients with tumors

with *MGMT* promoter methylation [54, 63]. The substantial toxicity profile limits combination with other agents.

3.4.2 Temozolomide monotherapy and combination regimens

Table 2 provides an overview on clinical trials of TMZ monotherapy, randomized trials with a TMZ monotherapy arm and combination regimens. TMZ was approved in 1999 for recurrent glioblastoma based on the data of 2 phase II trials.

TMZ was superior to procarbazine in patients, 60% of which were pretreated with nitrosoureas, with a PFS-6 rate of 21% vs. 8% and median OS prolonged by 1.5 months [46]. A single-arm phase II trial in 126 patients led to a PFS-6 rate of 18% [45]. Both trials used a schedule of TMZ 150-200 mg/m² for 5 out of 28 days. Other prospective studies mainly without previous TMZ treatment using this schedule showed similar PFS-6 rates ranging from 21 to 24% [64-67].

Alternative schedules of TMZ were developed aiming at overcoming TMZ resistance or exerting additional biologic effects, e.g. on angiogenesis. The main alternative schedules (Table 2) comprise low dose daily TMZ (40-50 mg/m²/d), 1-week-on/1-week-off (150 mg/m² for 7 days every 14 days), and 3-week-on/1-week-off (75-100 mg/m² for 21 days every 28 days). Grade 3 and grade 4 hematologic toxicities were observed in most of the studies without apparent differences between the schedules. Since most of the studies were conducted as single-arm monotherapy trials with different inclusion criteria e.g. regarding number of recurrences and pretreatment characteristics, comparison of efficacy outcome is limited. Yet, it seems very unlikely that there are relevant differences between the various dose-intensified TMZ regimens, and their superiority over standard dose TMZ for patients experiencing recurrence after a TMZ-free interval has not been demonstrated either. The RESCUE phase II trial evaluated the timing of TMZ rechallenge with prospective grouping of patients. Patients who received continuous TMZ for progression during the first 6 cycles of adjuvant TMZ and those patients with a treatment-free interval longer than 2 months showed comparable PFS-6 rates of 27.3 and 35.7% (median PFS 3.6 and 3.7 months) deriving more benefit than those patients receiving TMZ beyond the standard 6 adjuvant cycles but without a treatment-free interval (PFS-6 7.4%, median PFS 1.8 months) [68].

Beyond the randomized trial leading to the approval of TMZ mentioned above, two other randomized trials with TMZ monotherapy as one of the treatment arms have

been published, a direct comparison with the combination of procarbazine, lomustine and vincristine (PCV) before TMZ became first-line standard [69] and a negative trial for the epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor afatinib [70] (Table 2 and 4), indirectly supporting the role of TMZ at recurrence.

The DIRECTOR trial compared TMZ 120 mg/m² for 7 days every 14 days with 3-week-on/1-week-off 80 mg/m² for 21 days every 28 days and found no difference in outcome between these two regimens [71]. However, this trial established the role of *MGMT* promoter methylation as a prognostic marker for benefit of TMZ in recurrent glioblastoma patients, with a PFS-6 of 39.7% in patients with *MGMT* promoter-methylated tumors versus 6.9% without *MGMT* promoter methylation, in a prospective manner. The benefit of TMZ at recurrence thus seems limited to patients with tumors with *MGMT* promoter methylation.

Numerous studies evaluated TMZ-based combination regimens in recurrent glioblastoma. Beyond the combination with nitrosoureas already discussed, the following combination partners have been prospectively evaluated in single-arm designs:

The anti-vascular endothelial growth factor (VEGF) antibody bevacizumab [72-75], interferon- α 2b [76], the multikinase inhibitor sorafenib [77], O6-benzylguanine [78], irinotecan [79], cisplatin [80, 81], liposomal doxorubicine [82] and ABT-414, an EGFR-targeted antibody-drug conjugate conjugated to monomethylauristatin F [83]. So far, these trials have failed to provide convincing efficacy signals beyond single agent activity of TMZ.

3.4.3 Other classical non-alkylating chemotherapy

Beyond chemotherapy with alkylating agents (TMZ or nitrosoureas), other classical non-alkylating chemotherapeutics have been explored in the treatment of recurrent glioblastoma. A detailed listing of all trials with these agents, mostly conducted as single arm studies, would be beyond the scope of this review. Carboplatin (CABARET trial) and irinotecan (BRAIN trial) have been evaluated in randomized phase II trials as add-ons to bevacizumab (Table 3) [84, 85]. Both trials were negative showing no difference in outcome but additional toxicity in the treatment arms with carboplatin and irinotecan.

3.4.4 Bevacizumab monotherapy and combination regimens

Table 3 summarizes trials with single-arm monotherapy, randomized trials and single arm combination containing bevacizumab. Bevacizumab was approved in 2009 for the treatment of recurrent glioblastoma by the FDA and in several other countries based on the results of 2 uncontrolled phase II trials with overall response rates of about one third of the patients and with PFS-6 rates of 42.6 and 29% [85, 86]. In Europe, approval was refused because of the lack of a bevacizumab-free control arm. There are 9 phase II trials with a bevacizumab monotherapy arm, 3 of them single-arm studies [86-88], 5 trials compared bevacizumab with bevacizumab plus another agent, such as irinotecan [85], carboplatin [84], the histone-deacetylase inhibitor vorinostat [89], the multikinase inhibitor dasatinib [90] or lomustine [58]. Only one trial had a bevacizumab-free control arm with lomustine [54]. Most of these trials used bevacizumab at 10 mg/kg IV every 2 weeks except one study with 15 mg/kg IV every 3 weeks [87] and one study with 5 mg/kg IV every 3 weeks when combined with lomustine [58]. Toxicity of bevacizumab was mainly non-hematologic including hypertension, thromboembolic events, fatigue, and impaired wound healing. Overall response rates were similar in most trials with around one third of the patients mainly assessed by Macdonald criteria but also by RANO [54, 90], while one trial only reports 6%, assessed by RANO criteria [84]. PFS-6 rates ranged from 18% to 42.6% and median OS from 6.5 to 9.2 months. In the randomized trials adding another experimental agent to bevacizumab (irinotecan, carboplatin, vorinostat or dasatinib) versus bevacizumab alone, both arms showed comparable efficacy leading to the conclusion of poor efficacy of the experimental agent without valid evidence regarding the single agent activity of bevacizumab [84, 85, 89, 90]. The BELOB trial comprising a bevacizumab-free control arm showed comparable single agent activity of bevacizumab versus lomustine. However, as already discussed, median OS of the combination of bevacizumab and lomustine was superior to single agent activity in this phase II trial in contrast to the EORTC 26101 phase III trial showing no difference in OS of the combination versus lomustine alone [54, 56].

More than one dozen prospective trials evaluated bevacizumab in combination with other experimental agents in a single-arm design (Table 3). The trials combining bevacizumab with nitrosoureas or TMZ have already been mentioned in Table 1-2. Surprisingly, the combination of bevacizumab with TMZ [72-75] seems less promising than the combination with nitrosoureas [54, 58, 59]. However, no direct comparisons are available and the trials evaluating bevacizumab plus temozolomide were

conducted in part in highly pretreated patient populations [72-74]. Five phase II trials evaluated the combination of irinotecan with bevacizumab [91-95], two trials added a third combination partner, cetuximab [96] or carboplatin [97]. PFS-6 rates ranged between 28 and 50.3% and median OS between 6.7 and 9.7 months. Other experimental agents that were combined with bevacizumab are etoposide [98], the mTOR inhibitor temsirolimus [99], the EGFR-targeted tyrosine kinase inhibitor erlotinib [100], the multikinase inhibitor sorafenib [101], the histone deacetylase inhibitors panobinostat [102] or vorinostat [103].

No efficacy signal beyond the data for bevacizumab monotherapy was seen except for lomustine at phase II with negative results in the comparative phase III trial (EORTC 26101) [54, 56]; in addition, some of the combination trials reported increased toxicity.

In the clinic, after progression on a bevacizumab-containing regimen, bevacizumab is often continued fearing a rebound effect when stopping antiangiogenic therapy and data from metastatic colorectal cancer support this concept [104]. This clinical practice so far was lacking evidence from prospective and controlled clinical trials. This question was prospectively addressed with the cohort of the CABARET randomized phase II trial comparing bevacizumab monotherapy with bevacizumab plus carboplatin [105]. Upon progression of one of the two regimens, patients were randomized again to continue or cease bevacizumab. Patients previously receiving bevacizumab monotherapy could receive carboplatin or supportive care and those treated with combined bevacizumab plus carboplatin ceased carboplatin and could receive TMZ, etoposide or supportive care according to clinician choice. Forty-eight of initially 120 patients (40%) were available for randomization. There were no differences in outcome with a poor overall response of 0% in both arms, median PFS of the group continuing bevacizumab versus ceasing was 1.8 versus 2.0 months, OS was 3.4 versus 3.0 months.

In conclusion, there is little evidence to continue bevacizumab in patients with glioblastoma after progression on a bevacizumab-containing regimen.

3.4.5 Targeted therapy

In the last decade, a plethora of agents have been suggested for targeting essential pathways in glioblastoma. The pharmacologic agents had commonly been tested *in vitro* or *in vivo* or both before. Table 4 gives an overview on several phase II studies

conducted with agents targeting receptors or soluble factors involved in angiogenesis, oncogenic pathways or factors presumably involved in tumor cell stemness or tumor invasiveness. Most studies were designed as single-arm trials, some agents were tested in a randomized fashion against lomustine or TMZ. The results of single-arm phase II studies again are likely to be influenced by selection bias, patient heterogeneity and small patient numbers. Those agents tested against lomustine or TMZ failed to prove superiority.

Candidate anti-angiogenic agents other than bevacizumab, e.g. aflibercept, sunitinib, vandetanib or nintedanib failed to give an efficacy signal. Cediranib, a pan-tyrosine kinase inhibitor of VEGFR achieved similar response rates and PFS-6-data as did bevacizumab in phase II [85, 86, 106]. The subsequent phase III trial of cediranib showed comparable but not better efficacy than lomustine. Still, it is surprising that cediranib which is supposed to be similarly active as bevacizumab did not show superiority in combination with lomustine [52]. Further, the clinical development of a placental derived growth factor antibody was stopped early for presumed lack of activity [107].

Targeted agents aiming to inhibit oncogenic pathways such as enzastaurin, everolimus, PX-866, selinexor or dasatinib showed disappointing results [53, 108-111]. The phase II trial evaluating the multikinase inhibitor dasatinib only included patients with activation or overexpression of ≥ 2 putative dasatinib targets (ie, SRC, c-KIT, EPHA2, and PDGFR), nevertheless with poor efficacy results [111]. A placebo-controlled randomized phase II trial evaluating galunisertib, an inhibitor of transforming growth factor (TGF)- β receptor-1, shown to be involved in multiple pathogenic processes in glioblastoma *in vitro*, did not improve outcome when combined with lomustine [55]. However, the trial design containing a placebo-controlled active control (lomustine+placebo) at least allows reasonable conclusions on activity. The integrin inhibitor cilengitide tested in two dosing regimens in a randomized fashion, without a control arm lacking the experimental agent, showed a trend to prolonged survival at the higher dose [112]. A novel principle, the inhibition of histone deacetylase aiming to target epigenetic gene regulation, with agents such as vorinostat or panobinostat so far failed to provide efficacy signals [89, 102, 113, 114]. However, based on the mechanism, the combination with cytotoxic drugs such as lomustine or temozolomide might rather lead to synergistic effects than with bevacizumab. Several EGFR-targeting agents have been developed, such as

gefitinib, erlotinib or afatinib, a strategy effective in EGFR-mutated non-small-cell lung cancer, however, so far with disappointing results in glioblastoma [57, 70, 108, 115]. The phase II trial evaluating afatinib, an irreversible ErbB family blocker, showed that in the biomarker cohort (70 patients, 59%) the median PFS in those patients treated with afatinib monotherapy with EFGRvIII-positive tumors vs -negative tumors was 3.4 vs 1.0 months pointing towards efficacy in subgroups (over-)expressing the target [70]. The efficacy of ABT-414 might be further improved in a target-selected cohort since all patients of an unselected phase I cohort with confirmed response had EGFR amplification [83]. A current phase II trial (Clinicaltrials.gov NCT02343406) investigates whether ABT 414 alone or combined with temozolomide is superior to lomustine or temozolomide re-challenge in recurrent EGFR-amplified glioblastoma. In conclusion, so far none of the targeted agents has proven efficacy beyond the activity of alkylator therapy.

In conclusion, so far none of the targeted agents has proven efficacy beyond the activity of alkylator therapy.

3.4.6 Immunotherapeutic approaches

Therapeutic principles of immunotherapy include immunomodulatory drugs aiming at activating the immune system against the tumor, treatment with oncolytic viruses and different vaccination approaches, either cell-based or antigen-based or both. All approaches theoretically should work best if applied early in the course of the disease to patients with minimal residual disease. This is why the majority of immunotherapeutic studies in glioblastoma today are conducted in the first-line setting and no longer in recurrent glioblastoma.

A promising immunotherapeutic approach is “immune checkpoint inhibition” interfering with inhibitory T cell signaling via programmed death 1 (PD-1), the PD-1-ligand, or cytotoxic T-lymphocyte-associated antigen 4 (CTLA-4). Preliminary data of a phase II study, randomizing patients with recurrent glioblastoma to either nivolumab, a PD-1 antibody, alone or combined with ipilimumab, a CTLA-4 antibody, showed an overall survival rate of 75% at 6 months for both arms pooled [116].

Relevant toxicity including included colitis, cholecystitis, diabetic ketoacidosis, and confusion leading to discontinuation occurred in the combination arm in 50% of the patients. Other checkpoint inhibitors targeting PD-1 (Pembrolizumab, Clinicaltrials.gov NCT02337491) or PD-L1 (MEDI4736, Clinicaltrials.gov

NCT02336165) are currently evaluated in phase II.

A phase I trial in recurrent glioblastoma evaluated TMZ combined with a vaccine of monocyte-derived dendritic cells pulsed with autologous tumor cells previously been exposed to TMZ. Two of 9 patients had objective radiological responses and were progression-free at 6 months [117].

Gliovac, a vaccine composed of autologous and allogene antigens combined with GM-CSF and previous treatment with cyclophosphamide was shown to be safe in phase I and all patients (n=9) were alive at 6 months [118]. A phase II trial (n=41 patients) evaluated a vaccine composed of heat shock peptide protein complexes (HSPPC-96) after repeat tumor resection. PFS-6 of evaluable patients was 29.3%, median PFS 4.4 and median OS 9.8 months [119]. Strict patient selection after inclusion for being “evaluable” (30% were excluded) limits the interpretation of these results. A phase II trial showed promising results for rindopepimut, a vaccine consisting of a peptide sequence of EGFRvIII, plus bevacizumab versus bevacizumab plus control vaccine in bevacizumab-naïve patients (1st or 2nd relapse, n=72). The rindopepimut arm had higher overall response rate (24% vs. 17%), prolonged PFS-6 (27% vs. 11%) and median OS (12 vs. 8.8 months). An association of an anti-EGFRvIII immune titer generation with OS within the rindopepimut arm further supports this approach [120].

3.4.7 Other approaches

Beyond repeat surgery, irradiation, various chemotherapeutic and immunotherapeutic approaches, two further treatment concepts for recurrent glioblastoma shall be briefly mentioned. One approach comprises a portable device, called tumor-treating alternating electric fields (TTFields/NovoTTF), delivering low-intensity, intermediate frequency electric fields via non-invasive, transducer arrays aiming at physically interfere with cell division. NovoTTF(n=120 patients, 20-24h/d) was evaluated in a randomized phase III trial versus best physicians choice of chemotherapy (n=117 patients: bevacizumab in 31%, irinotecan in 31%, nitrosourea in 25%, carboplatin in 13%, temozolomide in 11% of patients). Median OS was comparable in both groups with 6.1 versus 6.6 months, PFS-6 was 21.4% and 15.1%. Toxicity was limited to moderate and mild skin reactions [121]. The interpretation of these data remains controversial, but the trial was overall as negative as many other trials discussed above.

Another “chemotherapy-free” approach represents the ERGO trial that examined a ketogenic diet in 20 patients with recurrent glioblastoma. With a median PFS of 1.2 and an OS of 7.4 months, there was no evidence of activity [122].

4. Summary and conclusion

Despite a plethora of clinical trials in recurrent glioblastoma, there are no established standards of care beyond alkylating chemotherapy or bevacizumab. The future role of bevacizumab is uncertain since the EORTC 26101 trial failed to demonstrate superiority for OS of lomustine plus bevacizumab over lomustine alone [56]. Many single-arm trials are difficult to interpret because of the lack of a controlled and randomized design and heterogeneous cohorts. Stratification of patients according to prognostic markers and implementation of control groups already in phase II trials may improve design and facilitate data interpretation in future trials. Based on the available clinical data for therapeutic strategies in glioblastoma, some conclusions, illustrated in Figure 1, can be drawn:

- The RANO criteria represent the currently most accepted approach for diagnosis of progression and response in recurrent glioblastoma and the international consensus MRI protocol provides a good working tool for practice and trials.
- The evidence for repeat surgery or reirradiation is limited, calling for prospective randomized trials determining their efficacy.
- Nitrosoureas still represent the most widely accepted standard option for systemic chemotherapy at recurrence
- *MGMT* promotor methylation may emerge as a predictive biomarker for benefit of TMZ rechallenge in recurrent glioblastoma, indirectly discouraging treatment with TMZ for *MGMT* promoter-unmethylated tumors. The best schedule of TMZ at recurrence has not been defined, and may be scheduling matters less than previously thought.
- There is clinical activity of bevacizumab monotherapy at recurrence, but an effect on overall survival is uncertain. Prospective data from phase II trials pointed towards efficacy of a combination regimen with nitrosoureas which was not confirmed in phase III.
- Immunotherapeutic concepts are currently under evaluation for newly diagnosed and recurrent glioblastoma, with promising preliminary data for rindopepimut, warranting further efforts exploring further immunotherapeutic agents even in the

recurrent setting, e.g., immune checkpoint inhibitors

In summary, treatment concepts for patients with recurrent glioblastoma in clinical practice should be somewhat individualized and take into account age, performance status, *MGMT* promoter methylation status and response to previous regimens as well as quality of life with regard to expected toxicities. To improve evidence on treatment options, patients should be treated within carefully designed clinical trials. Rebiopsies may be required more often to ascertain molecular tumor status as more targeted therapies become available. Further preclinical and clinical investigation is needed to improve the prognosis for patients with recurrent glioblastoma.

5. Conflicts of interest statement:

KS has received honoraria from Roche for advisory board participation.

WW has received research grants from Apogenix, Boehringer Ingelheim, Eli Lilly, immatics, MSD and Roche as well as honoraria for lectures or advisory board participation from MSD and Roche. WW is or has been the coordinating investigator for sponsored clinical trials evaluating APG101 (Apogenix), bevacizumab (Roche), galunisertib (Eli Lilly), temozolomide (MSD) and temsirolimus (Pfizer).

MW has received research grants from Acceleron, Actelion, Alpinia Institute, Bayer, Isarna, MSD, Merck & Co, Novocure, PIQUR and Roche and honoraria for lectures or advisory board participation or consulting from Celldex, Immunocellular Therapeutics, Isarna, Magforce, MSD, Merck & Co, Northwest Biotherapeutics, Novocure, Pfizer, Roche and Teva.

6. Tables

Table 1. Nitrosourea monotherapy and combination regimens¹

Design	No. of relapse, no. of patients	Pre-treatment	Treatment	ORR (%)	PFS-6 (%)	mPFS (months)	mOS (months)	Ref.
Single-arm monotherapy trials								
Phase II	n=40	RT	BCNU 80 mg/m ² on days 1-3 q8wk for max 6 cycles	15%	17.5%	NA	7.5	[47]
Phase II	1 st , n=27	RT/TMZ+TMZ	FOT IV 100 mg/m ² qwk for 3 wks (induction), then q3wk (maintenance)	30%	48.2%	5.7	9.1	[48]
Phase II	1 st , n=43	RT/TMZ+TMZ	FOT 75-100 mg/m ² qwk for 3 wks, after 5-wk break 100 mg/m ² q3wk for ≤1 yr	7%	20.9%	1.7	6.0	[49]
Phase II	1 st , n=50	RT/TMZ+TMZ	FOT 100 mg/m ² qwk for 3 wks (induction), 4- to 6-wk break, then FOT 100 mg/m ² q3wk (maintenance)	18%	52%	6.1	8.1	[51]
Phase II	1 st , n=40	RT/TMZ+TMZ	FOT IV 80 mg/m ² on days 1, 15, 30, 45, and 60 (induction), then 80 mg/m ² q4wk (maintenance)	25%	61%	6.1	11.1	[50]
Randomized trials								
Phase II, randomized	1 st , CTR: n=29 (BCNU), n=27 (TMZ), n=52 evaluable; n=54 (ERL)	RT+/-Chemo (64% CTR/ 65% ERL)	CTR (BCNU 60 mg/m ² on days 1-3 q8wk (max 5 cycles) or TMZ 150-200 mg/m ² (5/28) or ERL 150-200 mg/d	10% (CTR), 4% (ERL)	24% (CTR), 11% (ERL)	2.4 (CTR), 1.8 (ERL)	7.3 (CTR), 7.7 (ERL)	[57]
Phase III, randomized 2:1	1st: n=70 (CCNU), n=129 (Enzastaurin) 2nd: n=21 (CCNU), n=45 (Enzastaurin)	RT+/-Chemo (NA, no nitrosourea)	CCNU 100-130 mg/m ² q6wk or Enza 500 mg/d (1125 mg on day 1)	4% (CCNU), 3% (Enza)	19% (CCNU), 11% (Enza)	1.6 (CCNU), 1.5 (Enza)	7.1 (CCNU), 6.6 (Enza)	[53]
Phase III, double-blind, randomized 1:2:2	1 st , n=65 (CCNU) n=131 (CED), n=129 (CED+CCNU)	RT/TMZ+TMZ	CCNU 110 mg/m ² q6wk + placebo or CED 30 mg/d or CED 20 mg/d + CCNU 110 mg/m ² q6wk	8% (CCNU), 14 % (CED), 16%, (CED+CCNU)	24.5% (CCNU), 16% (CED), 34.5 % (CED+CCNU)	2.7 (CCNU), 3.1 (CED), 4.2 (CED+CCNU)	9.8 (CCNU), 8.0 (CED), 9.4 (CED+CCNU)	[52]
Phase II, randomized	1 st , n=50 (BEV), n=46 (CCNU), n=44 (BEV+CCNU)	RT/TMZ+TMZ	BEV 10 mg/kg q2wk or CCNU 110 mg/m ² q6wk or BEV q2wk +CCNU 90 (-110) mg/m ² q6wk	38% (BEV), 5% (CCNU), 34% (BEV+CCNU)	18% (BEV), 11% (CCNU), 41% (BEV+CCNU)	3 (BEV), 2 (CCNU), 11 (BEV+CCNU)	8 (BEV), 8 (CCNU), 11 (BEV+CCNU)	[54]
Phase II, randomized 2:1:1, placebo-controlled	NA, n=79 (CCNU+G), n=39 (G), n=40 (CCNU+P)	NA	CCNU q6w +Galunisertib (CCNU+G) or Galunisertib (300mg/d) for 14 days q28d alone (G)	1% (CCNU+G), 5% (G), 0% (CCNU+P)	NA	1.8 (CCNU+G), 1.8 (G), 1.9 (CCNU+P)	6.7 (CCNU+G), 8 (G), 7.5 (CCNU+P)	[55]

			or CCNU q6wk +placebo (P+CCNU)					
Phase II, randomized	1 st , 2 nd , 3 rd , n=68 evaluable pts	RT/TMZ+TMZ+/- other	BEV 10 mg/kg q2wk or BEV 5 mg/kg q3wk+ CCNU 90 mg/m ² q6wk	NA	NA	4.1 (BEV), 4.3 (BEV+CCNU)	NA (ITT); Pts with 1 st relapse: 8.8 (BEV), 13.1 (BEV+CCNU)	[58]
Phase III, randomized 2:1	1 st , n=288 (CCNU+BEV), n=149 (CCNU)	RT/TMZ+TMZ	CCNU 90 mg/m ² q6wk + BEV 10 mg/kg q2wk or CCNU 110 mg/m ² q6wk	NA	NA	4.2 (CCNU+BEV), 1.5 (CCNU)	9.1 (CCNU+BEV), 8.6 (CCNU)	[56]
Single-arm combination therapy trials								
Phase II	2 nd , n=42	RT+ TMZ	BCNU (100 mg/m ²) on days 1+CPT- 11 (175 mg/m ²) on days 1-3 q1wk for 4 wks, cycles of 6 wks	21.4%	30.3%	3.9	11.7	[60]
Phase II	NA, n=38	RT+/-chemo (10.5%)	BCNU (150 mg/m ²) +TMZ 550 mg/m ² q6wks	5.6%	21%	2.5	7.8	[61]
Prospective	NA, n=20 (n=10 evaluable for outcome)	RT/TMZ+TMZ	TMZ 150 mg/m ² for 7 days q14d +FOT 110 mg/m ² monthly q14d	NA	40%	4.3	NA	[62]
Phase II	1 st , n =54	RT/TMZ+TMZ	BEV 10 mg/kg q2wk +FOT 75 mg/m ² on day 1 and day 8, then q3wk	52%	42.6%	5.2	9.1	[59]

Abbreviations: BCNU, carmustine; CCNU, lomustine; CED, cediranib; CPT-11, irinotecan; CTR, control arm; d, day; Enza, enzastaurin; ERL, erlotinib; FOT, fotemustine; G: Galunisertib; ITT, intention-to-treat population; max, maximum; mOS, median overall survival; mPFS, median progression-free survival; NA, not available; P, placebo; PFS, progression-free survival; pts, patients; q, every; Ref, reference; RT, radiotherapy; TMZ, temozolomide; wk, week;

¹ All data presented for glioblastoma patients only. In case, the study was designed for different patient populations, only the respective data for glioblastoma patients were included in the table.

Table 2. Temozolomide monotherapy and combination regimens^{1/2}

Design	No. of relapse, no. of patients	Pre-treatment	Treatment	ORR (%)	PFS-6	mPFS (months)	mOS (months)	Ref.
Single-arm monotherapy trials								
Phase II	1 st , n=126	RT +/- nitrosourea (40%)	Chemo-naïve pts: TMZ 200 mg/m ² /d for 5 days q28d Pts with previous chemotherapy: TMZ 150 mg/m ² /d for 5 days q28d	8%	18%	2.1	5.4	[45]
Phase II	2 nd , n=22	RT+PCV	TMZ 150 mg/m ² /d for 5 days q28d	23%	32%	NA	7.6	[65]
Phase II	2 nd , n=42	RT+PCV	TMZ 150 mg/m ² /d for 5 days q28d	19%	24%	NA	7.0	[66]
Phase II	NA, n=28	RT (all but 1) +/- chemo (mostly nitrosourea)	TMZ 75 mg/m ² /d for 42 days q70d	0%	19%	2.3	7.7	[123]
Phase II	1 st , n=12; 2 nd and 3 rd , n=9	RT (all) +/- nitrosourea (43%)	TMZ 150 mg/m ² for 7 days q14d	10%	48%	4.9	NA	[124]
Prospective	1 st , n=13	RT	TMZ 200 mg/m ² /d for 5 days q28d, max 4 cycles	NA	21%	NA	NA	[67]
Phase II	1 st , n=33	RT	TMZ 75 mg/m ² /d for 21 days q28d	9%	30%	3.7	9.2	[125]
Prospective	NA, n=12	RT+TMZ+/-ACNU (n=1), PCV (n=2)	TMZ 40 mg/m ² /d	17%	NA	6	11	[126]
Prospective	NA, n=19	RT (all) +/- chemo (mostly nitrosourea)	TMZ 150-200 mg/m ² /d for 5 days q28d	24%	22%	2.2	9.9	[64]
Phase II	NA, n=64	RT+/-chemo (n=41 nitrosourea, n=9 TMZ+nitrosourea)	TMZ 150 mg/m ² for 7 days q14d	NA	43.8%	5.5	NA	[127]
Phase II	1 st , n=48; ≥2: n=20	RT+/- chemo (previous non-nitrosourea: chemo n=7; nitrosourea: n=33)	TMZ 200 mg/m ² initial dose, then 9 consecutive doses at 90-100 mg/m ² q12h for 28 days;	31%	35%	4.2	8.8	[128]
Phase II	NA, n=27 GB, n=20 AG	RT+chemo (93% TMZ 1 st line, 49% at 2 nd recurrence), some nitrosourea	TMZ 85 mg/m ² for 21 days q28d	7%	0	NA	5.1 ¹	[129]
Phase II	1 st , n=33 (early), n=27 (extended) n=28 (rechallenge) per timing of progression during/after adjuvant therapy	RT+TMZ/TMZ	50 mg/m ² /d continuous (max. 1 year)	NA	23.9% (all groups), 27.3% (early), 7.4% (extended), 35.7% (rechallenge)	NA (all groups) 3.6 (early), 1.8 (extended) 3.7 (rechallenge)	9.3 (all groups)	[68]
Phase II	1 st , n=38	RT+TMZ/TMZ	40-50 mg/m ² /d	5%	32.5%	3.9	9.4	[130]

Phase II	1 st , n=27	RT+TMZ/TMZ	TMZ 130 mg/m ² for 7 days q14d, max. 1 year	11.1%	33.3%	4.2	6.9	[131]
Phase II	1 st , n=58	RT+TMZ/TMZ	75-100 mg/m ² /d for 21 days q28d	13%	11%	1.8	11.7	[132]
Phase II	1 st , n=12, 2 nd , n=18, ≥3, n=10	RT+TMZ/TMZ+ other	TMZ 150 mg/m ² for 7 days q14d	NA	10%	1.8	5	[133]
Randomized trials								
Phase II, randomized	1 st , n=112 (TMZ), n=113 (PCB)	RT +/- nitrosourea (65% TMZ group, 68% PCB group)	TMZ 150-200 mg/m ² /d for 5 days q28d or PCB 125-150 mg/m ² /d for 28 days q56d	5% (TMZ), 5% (PCB)	21% (TMZ), 8% (PCB)	2.8 (TMZ), 1.9 (PCB)	NA	[46]
Phase II, randomized	1 st GBM and AA; n=87 (TMZ 5/28), n=81 (TMZ 21/28), n=162 (PCV)	RT	TMZ 200 mg/m ² for 5 days q28d (TMZ 5/28) or TMZ 100 mg/m ² for 21 days q28d (TMZ 21/28) or PCV	NA	NA	5.0 ² (TMZ 5/28), 4.2 ² (TMZ 21/28), 3.6 ² (PCV)	8.5 ² (TMZ 5/28), 6.6 ² (TMZ 21/28), 6.7 ² (PCV)	[69]
Phase II	NA, n=10 (TMZ+BEV), n=13 ETO+BEV)	RT+TMZ/TMZ +BEV+/-other	TMZ 50 mg/m ² /d + BEV 10 mg/kg q2wk or ETO 50 mg/m ² /d for 21 days q28d + BEV 10 mg/kg q2wk	0%	4.4%	1.0 (TMZ+BEV), 1.9 ETO+BEV)	NA	[74]
Phase II, randomized	1 st , n=39 (TMZ), n=41 (A), n=39 (A+TMZ)	RT+TMZ/TMZ	TMZ 75 mg/m ² /d for 21 days q28d (TMZ) or Afatinib 40mg/d (A) or Afatinib 40mg/d+ TMZ 75 mg/m ² /d for 21 days q28d (A+TMZ)	19% (TMZ) 2.4% (A), 5% (A+TMZ)	23% (TMZ) 3% (A) 10% (A+TMZ)	1.9 (TMZ) 1 (A) 1.5 (A+TMZ)	10.6 (TMZ) 9.8 (A) 8 (A+TMZ)	[70]
Phase II, randomized	1 st , n=105	RT+TMZ/TMZ	TMZ 120 mg/m ² for 7 days q14d (Arm A) or TMZ 80 mg/m ² /d for 21 days q28d (Arm B)	8% (Arm A) 16% (Arm B)	17.1% (Arm A) 25% (Arm B)	NA	9.8 (Arm A) 10.6 (Arm B)	[71]
Single-arm combination therapy trials								
The combination of TMZ+nitrosoureas is reported in Table1								
Phase II	1 st n=15 2 nd , n=15 3 rd , n=2	RT/TMZ+TMZ+/- other (BEV in n=4)	TMZ 50 mg/m ² /d + BEV 10 mg/kg q2wk	28%	18.8%	3.6	8.5	[72]
Phase I/II	NA, n=15	NA	TMZ 50 mg/m ² /d q3wk + BEV 10 mg/kg q3wk	NA	6.7%	2.4	3.6	[73]
Phase II	1 st , n=32	RT+TMZ/TMZ	TMZ 150 mg/m ² for 7 days q14d+ BEV 10 mg/kg q2wk	40.6%	21.9%	4.2	7.3	[75]
Phase II	NA, n=34 (TMZ+IFN), n=29 (TMZ+pIFN)	RT+/-chemo (no TMZ or IFN)	TMZ 150-200 mg/m ² /d for 5 days q28d + IFN-α2b: 4 MU/ m ² 3x/wk (TMZ+IFN) and TMZ 150-200 mg/m ² /d for 5 days q28d + Long-acting PEG-IFN-α2b 0.5 µg/kg/wk (TMZ+pIFN)	12% (TMZ+IFN) 3% (TMZ+pIFN)	31% (TMZ+IFN) 38% (TMZ+pIFN)	3.6 (TMZ+IFN) 4.4 (TMZ+pIFN)	7.2 (TMZ+IFN) 10 (TMZ+ pIFN)	[76]

Phase II	NA, 1-5 treatment lines, n=32	RT+/- chemo (TMZ, BEV and others)	TMZ 50 mg/m ² /d + Sorafenib 400 mg 2x/d	3%	9.4%	1.5	9.6	[77]
Phase II	NA, n=34	NA/TMZ (all)	TMZ 472 mg/m ² q28d +O(6)-BG 1-h infusion of 120 mg/m ² , followed immediately by a 48-h infusion of 30 mg/m ² q28d	3%	9	1.7	4.5	[78]
Phase I	NA, n=91	RT/chemo	TMZ 200 mg/m ² /d for 5 days q28d + CPT-11 40 mg/m ² to 375 mg/m ² IV on weeks 1, 2, 4, and 5 of each 6-wk cycle	12%	27.3%	2.6	NA	[79]
Phase II	1 st , n=22	RT+/- chemo (9%, no TMZ)	TMZ 200 mg/m ² for 5 days q28d + Liposomal DOX 40 mg/m ² IV on day 1 q4wk	18%	32%	3.6	8.2	[82]
Phase II	NA, n=20	RT+nitrosourea+ CIS	TMZ 200 mg/m ² on days 2-6 q4wk +CIS 40 mg/m ² , on days 1 and 2 q4wk	10%	35%	4.9	NA	[80]
Phase II	1 st , n=50	RT	TMZ 130 mg/m ² bolus followed by 9 doses of 70 mg/m ² q12h (total of 5 days) from day 2 q4wk, from 2 nd cycle: 100 mg/m ² mg/m ² q12h for 5 days q4wk +CIS 75 mg/m ² on day 1 q4wk	20.4%	34%	4.2	11.2	[81]
Phase I	NA, n=18	NA	ABT-414 0.5-1.5 mg/kg q14d + TMZ 150-200 mg/m ² for 5 days q28d	22.2%	NA	NA	NA	[83]

Abbreviations: (6)-BG: O6-benzylguanine; A, afatinib; AA, anaplastic astrocytoma; AG, anaplastic glioma; BEV, bevacizumab, CIS, cisplatin; d, day; DOX, doxorubicine; ETO, etoposide; GB, glioblastoma; IFN, interferon; mOS, median overall survival; mPFS, median progression-free survival; NA, not available; PCB, procarbazine; PCV, procarbazine, CCNU, and vincristine; PCB: procarbazine, PFS, progression-free survival; pts, patients; q, every; Ref, reference; RT, radiotherapy; TMZ, temozolomide. wk, week;

¹ Data presented for glioblastoma patients only. In case, the study was designed for different patient populations, only the respective data for glioblastoma patients were included in the table

² Reporting of data for glioblastoma patients only was not possible due to lack of reporting data separately in the studies of Berrocal and colleagues [129] with 27 patients with glioblastoma, 15 with anaplastic astrocytoma and 5 with miscellaneous brain tumors and the study of Brada and colleagues [69] with 277 patients with glioblastoma, 53 with anaplastic astrocytoma and 20 with miscellaneous brain tumors.

Table 3. Bevacizumab monotherapy and combination regimens^{1/2}

Design	No. of relapse, no. of patients	Pre-treatment	Regimen	ORR (%)	PFS-6 (%)	mPFS (months)	mOS (months)	Ref.
Single-arm monotherapy trials								
Phase II	1 st , n=48	RT/TMZ+ TMZ	BEV 10 mg/kg q2wk, addition of CPT-11 at progression	35%	29%	3.7	7.2	[86]
Phase II	1 st and other, n=50	RT/TMZ+ TMZ+ other	BEV 15 mg/kg q3wk	NA	25%	2.8	6.5	[87]
Phase II	1 st , n=29	RT/TMZ+ TMZ	BEV 10 mg/kg q2wk	27.6%	33.9%	3.3	10.5	[88]
Randomized trials								
Phase II, randomized	1 st , n=50 (BEV), n=46 (CCNU), n=44 (BEV+CCNU)	RT/TMZ+ TMZ	BEV 10 mg/kg q2wk or CCNU 110 mg/m ² q6wk or BEV q2wk +CCNU 90 (-110) mg/m ² q6wk	38 (BEV), 5 (CCNU), 34 (BEV+CCNU)	18 (BEV), 11 (CCNU), 41 (BEV+CCNU)	3 (BEV), 2 (CCNU), 11 (BEV+CCNU)	8 (BEV), 8 (CCNU), 11 (BEV+CCNU)	[54]
Phase II, randomized	1 st , 2 nd , 3 rd , n=68 evaluable pts	RT/TMZ+ TMZ+/- other	BEV 10 mg/kg q2wk or BEV 5 mg/kg q3wk+ CCNU 110 mg/m ² q6wk	NA	NA	4.1 (BEV), 4.3 (BEV+CCNU)	NA (ITT); Pts with 1 st relapse: 8.8 (BEV), 13.1 (BEV+CCNU)	[58]
Phase III, randomized 2:1	1 st , n=288 (CCNU+BEV), n=149 (CCNU)	RT/TMZ+TMZ	CCNU 90 mg/m ² q6wk + BEV 10 mg/kg q2wk or CCNU 110 mg/m ² q6wk	NA	NA	4.2 (CCNU+BEV), 1.5 (CCNU)	9.1 (CCNU+BEV), 8.6 (CCNU)	[56]
Phase II, randomized	1 st , n=69 (BEV) 2 nd , n=16 (BEV) 1 st , n=66 (BEV+CPT) 2 nd , n=16 (BEV+CPT)	RT/TMZ+ TMZ	BEV 10 mg/kg q2wk w/o CPT-11 340 mg/m2 or 125 mg/m2 q2wk	28 (BEV), 38 (BEV+CPT-11)	42.6 (BEV) 50.3 (BEV+CPT-11)	4.2 (BEV) 5.6 (BEV+CPT-11)	9.2 (BEV) 8.7 (BEV+CPT-11)	[85]
Phase II, randomized	1 st , n=55 (BEV), n=55 (BEV+Carbo)	RT/TMZ+ TMZ	BEV 10 mg/kg q2wk w/o Carbo (AUC 5) q4wk	6% (BEV), 14% (BEV+Carbo)	18% (BEV), 15% (BEV+Carbo)	3.5 (BEV), 3.5 (BEV+Carbo)	7.5 (BEV), 6.9 (BEV+Carbo)	[84]
Phase II, randomized	1 st , n=41 (BEV), n=49 (BEV+VOR)	RT/TMZ+ TMZ	BEV 10 mg/kg q2wk w/o VOR 400mg/d for 7 days q2wk	NA	NA	3.6 (BEV), 4.2 (BEV+VOR)	7.0 (BEV), 8.3 (BEV+VOR)	[89]
Phase II, randomized 1:2	NA, n=38 (BEV+P), n=83 (BEV+Dasa)	NA	BEV 10 mg/kg q2wk+P w/o Dasa 100 mg twice/d	26.5% (BEV+P), 18.3% (BEV+Dasa)	18.4% (BEV+P), 27.2% (BEV+Dasa)	NA	7.9 (BEV+P), 7.2 (BEV+Dasa)	[90]
Single-arm combination therapy trials								
The combination of BEV+nitrosoureas is reported in Table 1								
The combination of BEV+TMZ is reported in Table 2								
NA	NA, n=11 GB, n=10 other glioma	NA	BEV 5 mg/kg every other wk for 2 doses + CPT-11 125 mg/m ² qwk for 4 doses, followed by 2-wk rest period	43%	NA	NA	NA	[91]

Design	No. of relapse, no. of patients	Pre-treatment	Regimen	ORR (%)	PFS-6 (%)	mPFS (months)	mOS (months)	Ref.
Phase II	NA, n=23	RT and/or chemo	BEV 10 mg/kg q2wk+ CPT-11 340 mg/m ² (with EIAED) or 125 mg/m ² (without EIAED) q2wk	61%	30%	4.6	9.2	[93]
Phase II	NA, n=35	RT/TMZ+ TMZ+ other	<i>Cohort 1:</i> BEV 10 mg/kg q2wk + CPT-11 340 mg/m ² (with EIAED), or 125 mg/m ² (without EIAED), q2wk (6-wk cycle) <i>Cohort 2:</i> BEV 15 mg/kg IV q21d + CPT-11 340 mg/m ² (with EIAED) or 125 mg/m ² (without EIAED) on days 1, 8, 22, and 29 (6-wk cycle)	NA	46% ³	5.5 ³	9.7 ³	[92]
Phase II	NA, n=57	RT/TMZ+TMZ	BEV 10 mg/kg +CPT-11 200 mg/m ² q2wk	NA	37%	NA	NA	[94]
Phase II	NA, n=32	NA	BEV 10 mg/kg q2wk + CPT-11 340 mg/m ² IV (with EIAED) 125 mg/m ² (without EIAED) q2wk	25%	28%	5.2	7.9	[95]
Phase II	1 st within last 6 months of first-line treatment, n=43 (n=32 evaluable)	RT/TMZ+ TMZ	BEV 5-10 mg/kg q2wk + CPT-11 340 mg/m ² IV (with EIAED) 125 mg/m ² (without EIAED) q2wk + CET 400 mg/m ² IV as loading dose, followed by 250 mg/m ² /wk	34%	30%	3.7	6.7	[96]
Phase II	1 st , n=27, 2 nd , n=11, 3 rd , n=2	RT/TMZ+ TMZ+ other (no BEV)	BEV 10 mg/kg q2wk + CPT-11 340 mg/m ² IV (with EIAED) 125 mg/m ² (without EIAED) q2wk + Carbo (AUC 5) q4wk	33%	46.5%	5.9	8.3	[97]
Phase II	1 st , n=14, 2 nd , n=8, 3 rd , n=5	RT+chemo	BEV 10 mg/kg q2w + ETO 50 mg/m ² daily for 21 consecutive days each month	22%	44.4%	4.1	10.2	[98]
Phase II	1st: 52%; 2nd: 36%; 3rd: 12%	RT/TMZ+ TMZ	BEV 10 mg/kg q2wk + ERL 500 mg/kg/d (with EIAED) or, 200 mg/kg/d (without EIAED)	50%	29.2%	4.1	10.3	[100]
Phase II	NA, n=13	RT/TMZ+ TMZ+BEV	BEV 10 mg/kg q2wk starting on day 8 + temsirolimus 25 mg qwk starting on day 1	0%	NA	1.8	3.5	[99]
Phase II	NA, n=54	RT/TMZ+ TMZ+/- other	BEV 5 mg/kg q2wk + Sorafenib 100-200 mg/d	18.5%	20.4%	2.9	5.6	[101]
Phase II	≤2 relapses, n=24 (interim analysis)	NA (No anti-VEGF, no HDAC-I)	BEV 10 mg/kg q2wk +Panobinostat 30mg 3 times/wk	29.2%	30.4%	5	9	[102]
Phase II	NA, n=40	NA	BEV 10 mg/kg q2wk +VOR 400mg/d for 7 days q2wk	22.5%	30%	NA	10.4	[103]

Abbreviations: AUC, area under the curve; BEV, bevacizumab; CET, cetuximab; Carbo, carboplatin; CPT-11, irinotecan; d, day; Dasa: Dasatinib; EIAED, enzyme-inducing antiepileptic drugs; ERL, erlotinib; ETO, etoposide; GB, glioblastoma; HDAC-I: histone deacetylase-inhibitors, ITT: intention-to-treat-population; mOS, median overall survival; mPFS, median progression-free survival; NA, not available; P: placebo; PFS, progression-free survival; pts, patients; q, every; Ref, reference; RT, radiotherapy; TMZ, temozolomide,; VEGF: vascular endothelial growth factor, VOR: vorinostat; w/o, with or without; wk, week;

¹ Data presented for glioblastoma patients only. In case, the study was designed for different patient populations, only the respective data for glioblastoma patients were included in the table

² Reporting of data for glioblastoma patients only was not possible due to lack of reporting data for separately for one study [91] where combined data of patients were presented (11 patients with glioblastoma and 10 patients with anaplastic glioma)

³ Data presented for both cohorts combined

Table 4: Targeted agents, monotherapy and combination regimens^{1/2}

Design	No. of relapse, no. of patients	Pre-treatment	Regimen	Target	ORR	PFS-6	mPFS (months)	mOS (months)	Ref.
Phase II	NA, n=20	RT+chemo	Gefitinib 250 qd and everolimus 70 mg qw	EGFR (Gefitinib), mTOR (Everolimus)	14%	NA	2.6	5.8	[108]
Phase II	NA, n=31	RT or RT/TMZ+ TMZ	Cediranib 45 mg/d	VEGFR1-3, c-Kit, PDGFR	27%	25.8	3.8	7.5	[106]
Phase II	NA/34% 2 nd , total: n=78 (125mg), n=46 (175mg)	NA/Previous AAT (n=42)	XL184 125 mg <i>or</i> 175 mg qd	c-MET, VEGFR2, RET	NA	NA (125 mg) 21% (175 mg)	NA	NA	[134, 135]
Phase II	NA, n=35	NA	Pazopanib 800 mg qd	VEGFR1-3, c-Kit, PDGFR	6%	3%	2.8	NA	[136]
Phase II	1 st , n=37	RT+chemo	Sagopilone 16 mg/m ² q21d	Microtubule	0%	6.7%	1.4	7.6	[137]
Phase II	1 st n=39 2 nd , n=3 (excluded from efficacy analysis)	RT/TMZ+ TMZ	Aflibercept 4 mg/kg q2wk	VEGF, PIGF	18%	7.7	2.8	9.0	[138]
Phase II	NA, n=66	NA	VOR 400mg/d for 14 days of a 21-day cycle	Histone deacetylase	3%	15.2%	1.9	5.7	[113]
Phase II	≤3 relapse, n=16	RT/TMZ+ TMZ+/-other (BEV 48%)	AMG-102: 10 or 20mg/kg q2k	HGF	2%	NA	1.0 (10 mg) 1.2 (20 mg)	6.5 (10 mg) 5.4 (20 mg)	[139]
Phase II	1 st (n=18), 2 nd (n=7), 3 rd or more (n=7)	RT+chemo+/-others	Vandetanib 300 mg qd	VEGFR2, EGFR, "rearranged-during-transfection oncogene"	13%	6.5%	1.3	6.3	[140]
Phase II	≤2 relapse, n=16	RT/TMZ+ TMZ+/-other	Sunitinib 50mg/d for 4 wk every 6 wks	VEGFR1-2, c-Kit, PDGFR, FLT3, CSF-1R, RET	0%	16.7%	1.4	12.6	[141]
Phase II	NA, n=37	NA	VOR 400mg/d for 14 days of a 21-day cycle Bortezomib 1.3 mg/m ² on days 1, 4, 8, and 11	Histone deacetylase (VOR), Proteasom (Bortezomib)	3%	0%	1.5	3.2	[114]
Phase II	NA, n=56	RT+chemo	Erlotinib 150 mg/d, sorafenib 400mg twice/d	EGFR (Erlotinib), VEGFR, PDGFR, Raf kinases	5%	14%	2.5	5.7	[115]
Phase I	NA, n=18	NA	ABT-414 0.5-1.5 mg/kg q14d + TMZ 150-200 mg/m ² for 5 days q28d	EGFR (ABT-414)	22.2%	NA	NA	NA	[83]
Phase II	1 st , n=13 2 nd , n=12	RT/TMZ+ TMZ (n=13), BEV at 1 st relapse (n=12)	Nintedanib 200 mg bid	VEGFR1-3, FGFR1-3, PDGFR	0%	NA	1	6	[142]
Phase II	NA, n=41	RT+ chemo, others	Pazopanib 400 mg qd, Lapatinib 1000 mg ad	VEGFR1-3, c-Kit, PDGFR (Pazopanib), EGFR (Lapatinib)	5%	7.5%	1.9	NA	[143]
Phase II	NA, n=32 BEV naïve, n=31 BEV resistant	RT/TMZ+ TMZ +/- BEV or other	Sunitinib 37.5 mg qd	VEGFR1-2, c-Kit, PDGFR, FLT3, CSF-1R, RET	10% (BEV-naïve) 0% (BEV-resistant)	10.4% (BEV-naïve) 0% (BEV-resistant)	1 (BEV-naïve) 1 (BEV-resistant)	9.4 (BEV-naïve), 4.4 (BEV-resistant)	[144]
Phase II	≤2 relapse, n=31 (BEV-naïve), n=25 (previous BEV)	RT/TMZ+ TMZ+/-other	Verubulin 3.3 mg/m ² qwk for 3 wks, every 4 wks	Microtubule destabilizing, vascular disruption	10%(BEV-naïve), 4.2% (previous BEV)	14%(BEV-naïve), 8% (previous BEV)	1.6 (BEV-naïve), 0.8 (previous BEV)	9.4 (BEV-naïve), 3.4 (previous BEV)	[145]

Phase II	1 st , n=50, overexpression of at least 2 putative dasatinib targets	RT/TMZ+ TMZ	Dasatinib 100 mg -200 twice/d,	SRC, c-KIT, EPHA2, PDGFR	0%	6%	1.7	7.9	[111]
Phase II	NA, n=7 (Arm A), n=15 (Arm B, 12 evaluable)	RT/TMZ+ TMZ	3 doses of Selinexor 50 mg/m ² prior to surgery, Selinexor 50 mg/m ² after surgery (Arm A) <i>and</i> Selinexor 50 mg/m ² q2wk (Arm B)	Nuclear retention via XPO1 of p53, pRB, CDKN2A, p21 and FOXO	17%	NA	NA	NA	[109]
Phase II	1 st , n=33	RT/TMZ+ TMZ	PX-866 8mg/d	PI3K	3%	17%	NA	NA	[110]
Randomized trials									
Phase II, randomized	1 st , n=41 (500mg) n=40 (200mg)	RT, TMZ and others	CIL 500 <i>or</i> 2000 mg, 2x/wk	avβ3/avβ5 integrins	5% (500 mg) 13% (2000 mg)	10% (500 mg) ² 15% (2000 mg) ²	NA	6.5 ² (500 mg) 9.9 ² (2000 mg)	[112]
Phase II, randomized	1 st , CTR: n=29 (BCNU), n=27 (TMZ), n=52 evaluable; n=54 (ERL)	RT+/-Chemo (64% CTR/ 65% ERL)	CTR (BCNU 60 mg/m ² on days 1-3 q8wk (max 5 cycles) <i>or</i> TMZ 150-200 mg/m ² (5/28) <i>or</i> ERL 150-200 mg/d	EGFR (ERL)	10% (CTR), 4% (ERL)	24% (CTR), 11% (ERL)	2.4 (CTR), 1.8 (ERL)	7.3 (CTR), 7.7 (ERL)	[57]
Phase II, randomized	1 st and 2 nd , total n=26	RT+ chemo	CIL 500 <i>or</i> 2000 mg × 3 doses, resection, then 2000 mg 2x/wk	avβ3/avβ5 integrins	NA	12% (both arms)	1.9 (both arms)	NA	[146]
Phase III, double-blind, randomized 1:2:2	1 st , n=65 (CCNU) n=131 (CED), n=129 (CED+CCNU)	RT/TMZ+ TMZ	CCNU 110 mg/m ² q6wk + placebo <i>or</i> CED 30 mg/d <i>or</i> CED 20 mg/d + CCNU 110 mg/m ² q6wk	VEGFR1-3 and PDGFR (CED)	8% (CCNU), 14 % (CED), 16%, (CED+ CCNU)	24.5% (CCNU), 16% (CED), 34.5 % (CED+ CCNU)	2.7 (C2CCNU), 3.1 (CED11), 4.2 (CED+ CCNU)	9.8 (CCNU), 8.0 (CED), 9.4 (CED+ CCNU)	[52]
Phase III, randomized 2:1	1st: n=70 (CCNU), n=129 (Enzastaurin) 2nd: n=21 (CCNU), n=45 (Enzastaurin)	RT+/-Chemo (NA, no nitrosourea)	CCNU 100-130 mg/m ² q6wk <i>or</i> Enza 500 mg/d (1125 mg on day 1)	Protein kinase C, PI3K, AKT	4% (CCNU), 3% (Enza)	19% (CCNU), 11% (Enza)	1.6 (CCNU), 1.5 (Enza)	7.1 (CCNU), 6.6 (Enza)	[53]
Phase II, randomized 2:1:1, placebo-controlled	NA, n=79 (CCNU+G), n=39 (G), n=40 (CCNU+P)	NA	CCNU q6w + Galunisertib (CCNU+G) <i>or</i> Galunisertib (300mg/d) for 14 days q28d alone (G) <i>or</i> CCNU q6w +P (P+CCNU)	TGFβR1 (G)	1% (CCNU+G), 5% (G), 0% (CCNU+P)	NA	1.8 (CCNU+G), 1.8 (G), 1.9 (CCNU+P)	6.7 (CCNU+G), 8 (G), 7.5 (CCNU+P)	[55]
Phase II, randomized	1 st , n=39 (TMZ), n=41 (A), n=39 (A+TMZ)	RT/TMZ+ TMZ	TMZ 75 mg/m ² /d for 21 days q28d (TMZ) <i>or</i> Afatinib 40mg/d (A) <i>or</i> Afatinib 40mg/d+ TMZ 75	ErbB, EGFR (A)	19% (TMZ) 2.4% (A), 5% (A+TMZ)	23% (TMZ) 3% (A) 10% (A+TMZ)	1.9 (TMZ) 1 (A) 1.5 (A+TMZ)	10.6 (TMZ) 9.8 (A) 8 (A+TMZ)	[70]

			mg/m ² /d for 21 days q28d (A+TMZ)						
Phase II, randomized	1 st , n=41 (BEV), n=49 (BEV+VOR)	RT/TMZ+ TMZ	BEV 10 mg/kg q2wk w/o VOR 400mg/d for 7 days q2wk	Histone deacetylase	NA	NA	3.6 (BEV), 4.2 (BEV+VOR)	7.0 (BEV), 8.3 (BEV+VOR)	[89]
Phase II, randomized 1:2	NA, n=38 (BEV+P), n=83 (BEV+Dasa)	NA	BEV 10 mg/kg q2wk+P w/o Dasa 100 mg twice/d	SRC, c-KIT, EPHA2, PDGFR (Dasa)	26.5% (BEV+P), 18.3% (BEV+Dasa)	18.4% (BEV+P), 27.2% (BEV+Dasa)	NA	7.9 (BEV+P), 7.2 (BEV+Dasa)	[90]

Abbreviations: A, Afatinib; AAT, antiangiogenic therapy; BEV, bevacizumab; CCNU, lomustine; CED, cediranib; CIL, cilengitide; CSF-1R, Colony stimulating factor 1 receptor; d, day; Dasa: Dasatinib; EGFR, epidermal growth factor receptor; EPHA2, ephrin type-A receptor 2; ErbB, erythroblastic leukemia viral oncogene.; ERL, erlotinib; FGFR, fibroblast growth factor receptor; G: galunisertib; HGF, hepatocyte growth factor/scatter factor; c-MET, hepatocyte growth factor receptor; mOS, median overall survival; mPFS, median progression-free survival; mTOR, mammalian target of rapamycin; NA, not available; P, placebo; PDGF-R, platelet-derived growth factor receptor; PFS, progression-free survival; PI3K, phosphatidylinositol 3-kinase; PIGF, placental growth factor; pts, patients; q, every; Ref, reference; TMZ, temozolomide; TGFβR1, transforming growth factor-β-receptor-1; VEGF, vascular endothelial growth factor VEGFR, receptor to vascular endothelial growth factor; VOR, vorinostat; FLT-3, fms-like tyrosine kinase 3; w/o, with or without; wk, week;

¹ Data presented for glioblastoma patients only. In case, the study was designed for different patient populations, only the respective data for glioblastoma patients were included in the table

² Reporting of data for glioblastoma patients only was not possible due to lack of reporting data separately for one study where central histopathology review showed that 6 patients had either anaplastic astrocytoma or low-grade glioma [112]

7. Vitae

Katharina Seystahl, MD works as a Physician-Scientist at the Department of Neurology at the University Hospital Zurich. Research interests include clinical and molecular neuro-oncology with focus on antiangiogenic therapies and involvement of TGF- β in the pathogenesis of glioblastoma.

Wolfgang Wick, MD, is Chairman of the Neurology Clinic at the University Hospital Heidelberg, Germany and Division Head at the German Cancer Research Center. His research focus is on precision medicine and immunotherapies for brain tumor patients. He chairs the Neurooncology Working Group (NOA) of the German Cancer Society.

Michael Weller, MD is Chairman of the Department of Neurology at the University Hospital Zurich. His research interests include brain tumor immunology and clinical trials in neuro-oncology. He serves as the Chairman of the Brain Tumor Group of the European Organisation for Research and Treatment of Cancer (EORTC).

8. References:

1. Johnson, D.R. and O'Neill, B.P., Glioblastoma survival in the United States before and during the temozolomide era. *J Neurooncol* 2012, 107: 359-64.
2. Stupp, R., Mason, W.P., van den Bent, M.J., Weller, M., Fisher, B., Taphoorn, M.J., et al., Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med* 2005, 352: 987-96.
3. Hegi, M.E., Diserens, A.C., Gorlia, T., Hamou, M.F., de Tribolet, N., Weller, M., et al., MGMT gene silencing and benefit from temozolomide in glioblastoma. *N Engl J Med* 2005, 352: 997-1003.
4. Keime-Guibert, F., Chinot, O., Taillandier, L., Cartalat-Carel, S., Frenay, M., Kantor, G., et al., Radiotherapy for glioblastoma in the elderly. *N Engl J Med* 2007, 356: 1527-35.
5. Wick, W., Platten, M., Meisner, C., Felsberg, J., Tabatabai, G., Simon, M., et al., Temozolomide chemotherapy alone versus radiotherapy alone for malignant astrocytoma in the elderly: the NOA-08 randomised, phase 3 trial. *Lancet Oncol* 2012, 13: 707-15.
6. Malmstrom, A., Gronberg, B.H., Marosi, C., Stupp, R., Frappaz, D., Schultz, H., et al., Temozolomide versus standard 6-week radiotherapy versus hypofractionated radiotherapy in patients older than 60 years with glioblastoma: the Nordic randomised, phase 3 trial. *Lancet Oncol* 2012, 13: 916-26.
7. Weller, M., van den Bent, M., Hopkins, K., Tonn, J.C., Stupp, R., Falini, A., et al., EANO guideline for the diagnosis and treatment of anaplastic gliomas and glioblastoma. *Lancet Oncol* 2014, 15: e395-403.
8. Weller, M., Cloughesy, T., Perry, J.R., and Wick, W., Standards of care for treatment of recurrent glioblastoma--are we there yet? *Neuro Oncol* 2013, 15: 4-27.
9. Macdonald, D.R., Cascino, T.L., Schold, S.C., Jr., and Cairncross, J.G., Response criteria for phase II studies of supratentorial malignant glioma. *J Clin Oncol* 1990, 8: 1277-80.
10. Wen, P.Y., Macdonald, D.R., Reardon, D.A., Cloughesy, T.F., Sorensen, A.G., Galanis, E., et al., Updated response assessment criteria for high-grade gliomas: response assessment in neuro-oncology working group. *J Clin Oncol* 2010, 28: 1963-72.
11. Ellingson, B.M., Bendszus, M., Boxerman, J., Barboriak, D., Erickson, B.J., Smits, M., et al., Consensus recommendations for a standardized Brain Tumor Imaging Protocol in clinical trials. *Neuro Oncol* 2015, 17: 1188-98.
12. Kickingeder, P., Wiestler, B., Burth, S., Wick, A., Nowosielski, M., Heiland, S., et al., Relative cerebral blood volume is a potential predictive imaging

- biomarker of bevacizumab efficacy in recurrent glioblastoma. *Neuro Oncol* 2015, 17: 1139-47.
13. Ratai, E.M., Zhang, Z., Snyder, B.S., Boxerman, J.L., Safriel, Y., McKinstry, R.C., et al., Magnetic resonance spectroscopy as an early indicator of response to anti-angiogenic therapy in patients with recurrent glioblastoma: RTOG 0625/ACRIN 6677. *Neuro Oncol* 2013, 15: 936-44.
 14. Schmainda, K.M., Prah, M., Connelly, J., Rand, S.D., Hoffman, R.G., Mueller, W., et al., Dynamic-susceptibility contrast agent MRI measures of relative cerebral blood volume predict response to bevacizumab in recurrent high-grade glioma. *Neuro Oncol* 2014, 16: 880-8.
 15. Ellingson, B.M., Sahebjam, S., Kim, H.J., Pope, W.B., Harris, R.J., Woodworth, D.C., et al., Pretreatment ADC histogram analysis is a predictive imaging biomarker for bevacizumab treatment but not chemotherapy in recurrent glioblastoma. *AJNR Am J Neuroradiol* 2014, 35: 673-9.
 16. Kickingeder, P., Wiestler, B., Graf, M., Heiland, S., Schlemmer, H.P., Wick, W., et al., Evaluation of dynamic contrast-enhanced MRI derived microvascular permeability in recurrent glioblastoma treated with bevacizumab. *J Neurooncol* 2015, 121: 373-80.
 17. Radbruch, A., Eidel, O., Wiestler, B., Paech, D., Burth, S., Kickingeder, P., et al., Quantification of tumor vessels in glioblastoma patients using time-of-flight angiography at 7 Tesla: a feasibility study. *PLoS One* 2014, 9: e110727.
 18. Bonekamp, D., Deike, K., Wiestler, B., Wick, W., Bendszus, M., Radbruch, A., et al., Association of overall survival in patients with newly diagnosed glioblastoma with contrast-enhanced perfusion MRI: Comparison of intraindividually matched T1 - and T2 (*) -based bolus techniques. *J Magn Reson Imaging* 2015, 42: 87-96.
 19. Galldiks, N., Dunkl, V., Stoffels, G., Hutterer, M., Rapp, M., Sabel, M., et al., Diagnosis of pseudoprogression in patients with glioblastoma using O-(2-[¹⁸F]fluoroethyl)-L-tyrosine PET. *Eur J Nucl Med Mol Imaging* 2015, 42: 685-95.
 20. Guyotat, J., Signorelli, F., Frappaz, D., Madarassy, G., Ricci, A.C., and Bret, P., Is reoperation for recurrence of glioblastoma justified? *Oncol Rep* 2000, 7: 899-904.
 21. Bloch, O., Han, S.J., Cha, S., Sun, M.Z., Aghi, M.K., McDermott, M.W., et al., Impact of extent of resection for recurrent glioblastoma on overall survival: clinical article. *J Neurosurg* 2012, 117: 1032-8.
 22. McNamara, M.G., Lwin, Z., Jiang, H., Templeton, A.J., Zadeh, G., Bernstein, M., et al., Factors impacting survival following second surgery in patients with glioblastoma in the temozolomide treatment era, incorporating neutrophil/lymphocyte ratio and time to first progression. *J Neurooncol* 2014, 117: 147-52.

23. Woernle, C.M., Peus, D., Hofer, S., Rushing, E.J., Held, U., Bozinov, O., et al., Efficacy of Surgery and Further Treatment of Progressive Glioblastoma. *World Neurosurg* 2015.
24. Franceschi, E., Bartolotti, M., Tosoni, A., Bartolini, S., Sturiale, C., Fioravanti, A., et al., The effect of re-operation on survival in patients with recurrent glioblastoma. *Anticancer Res* 2015, 35: 1743-8.
25. Barbagallo, G.M., Jenkinson, M.D., and Brodbelt, A.R., 'Recurrent' glioblastoma multiforme, when should we reoperate? *Br J Neurosurg* 2008, 22: 452-5.
26. Park, J.K., Hodges, T., Arko, L., Shen, M., Dello Iacono, D., McNabb, A., et al., Scale to predict survival after surgery for recurrent glioblastoma multiforme. *J Clin Oncol* 2010, 28: 3838-43.
27. Park, C.K., Kim, J.H., Nam, D.H., Kim, C.Y., Chung, S.B., Kim, Y.H., et al., A practical scoring system to determine whether to proceed with surgical resection in recurrent glioblastoma. *Neuro Oncol* 2013, 15: 1096-101.
28. Nava, F., Tramacere, I., Fittipaldo, A., Bruzzzone, M.G., Dimeco, F., Fariselli, L., et al., Survival effect of first- and second-line treatments for patients with primary glioblastoma: a cohort study from a prospective registry, 1997-2010. *Neuro Oncol* 2014, 16: 719-27.
29. Gorlia, T., Stupp, R., Brandes, A.A., Rampling, R.R., Fumoleau, P., Ditttrich, C., et al., New prognostic factors and calculators for outcome prediction in patients with recurrent glioblastoma: a pooled analysis of EORTC Brain Tumour Group phase I and II clinical trials. *Eur J Cancer* 2012, 48: 1176-84.
30. Suchorska, B., Weller, M., Tabatabai, G., Senft, C., Hau, P., Sabel, M., et al., Complete Resection of Contrast-Enhancing Tumor Volume is Associated With Improved Survival in Recurrent Glioblastoma Results From the DIRECTOR Trial. *Neurosurgery* 2015, 62 Suppl 1: 209.
31. Yong, R.L., Wu, T., Mihatov, N., Shen, M.J., Brown, M.A., Zaghloul, K.A., et al., Residual tumor volume and patient survival following reoperation for recurrent glioblastoma. *J Neurosurg* 2014, 121: 802-9.
32. Ryu, S., Buatti, J.M., Morris, A., Kalkanis, S.N., Ryken, T.C., Olson, J.J., et al., The role of radiotherapy in the management of progressive glioblastoma : a systematic review and evidence-based clinical practice guideline. *J Neurooncol* 2014, 118: 489-99.
33. Combs, S.E., Debus, J., and Schulz-Ertner, D., Radiotherapeutic alternatives for previously irradiated recurrent gliomas. *BMC Cancer* 2007, 7: 167.
34. Dhermain, F., de Crevoisier, R., Parker, F., Gioloca, C., Kaliski, A., Beaudre, A., et al., [Role of radiotherapy in recurrent gliomas]. *Bull Cancer* 2004, 91: 883-9.

35. Fogh, S.E., Andrews, D.W., Glass, J., Curran, W., Glass, C., Champ, C., et al., Hypofractionated stereotactic radiation therapy: an effective therapy for recurrent high-grade gliomas. *J Clin Oncol* 2010, 28: 3048-53.
36. Miwa, K., Matsuo, M., Ogawa, S., Shinoda, J., Yokoyama, K., Yamada, J., et al., Re-irradiation of recurrent glioblastoma multiforme using ¹¹C-methionine PET/CT/MRI image fusion for hypofractionated stereotactic radiotherapy by intensity modulated radiation therapy. *Radiat Oncol* 2014, 9: 181.
37. Combs, S.E., Edler, L., Rausch, R., Welzel, T., Wick, W., and Debus, J., Generation and validation of a prognostic score to predict outcome after re-irradiation of recurrent glioma. *Acta Oncol* 2013, 52: 147-52.
38. Grosu, A.L., Weber, W.A., Franz, M., Stark, S., Piert, M., Thamm, R., et al., Reirradiation of recurrent high-grade gliomas using amino acid PET (SPECT)/CT/MRI image fusion to determine gross tumor volume for stereotactic fractionated radiotherapy. *Int J Radiat Oncol Biol Phys* 2005, 63: 511-9.
39. Wick, W., Fricke, H., Junge, K., Kobayakov, G., Martens, T., Heese, O., et al., A phase II, randomized, study of weekly APG101+reirradiation versus reirradiation in progressive glioblastoma. *Clin Cancer Res* 2014, 20: 6304-13.
40. Hundsberger, T., Brugge, D., Putora, P.M., Weder, P., Weber, J., and Plasswilm, L., Re-irradiation with and without bevacizumab as salvage therapy for recurrent or progressive high-grade gliomas. *J Neurooncol* 2013, 112: 133-9.
41. Flieger, M., Ganswindt, U., Schwarz, S.B., Kreth, F.W., Tonn, J.C., la Fougere, C., et al., Re-irradiation and bevacizumab in recurrent high-grade glioma: an effective treatment option. *J Neurooncol* 2014, 117: 337-45.
42. Minniti, G., Agolli, L., Falco, T., Scaringi, C., Lanzetta, G., Caporello, P., et al., Hypofractionated stereotactic radiotherapy in combination with bevacizumab or fotemustine for patients with progressive malignant gliomas. *J Neurooncol* 2015, 122: 559-66.
43. Cabrera, A.R., Cuneo, K.C., Desjardins, A., Sampson, J.H., McSherry, F., Herndon, J.E., 2nd, et al., Concurrent stereotactic radiosurgery and bevacizumab in recurrent malignant gliomas: a prospective trial. *Int J Radiat Oncol Biol Phys* 2013, 86: 873-9.
44. Lamborn, K.R., Yung, W.K., Chang, S.M., Wen, P.Y., Cloughesy, T.F., DeAngelis, L.M., et al., Progression-free survival: an important end point in evaluating therapy for recurrent high-grade gliomas. *Neuro Oncol* 2008, 10: 162-70.
45. Brada, M., Hoang-Xuan, K., Rampling, R., Dietrich, P.Y., Dirix, L.Y., Macdonald, D., et al., Multicenter phase II trial of temozolomide in patients with glioblastoma multiforme at first relapse. *Ann Oncol* 2001, 12: 259-66.

46. Yung, W.K., Albright, R.E., Olson, J., Fredericks, R., Fink, K., Prados, M.D., et al., A phase II study of temozolomide vs. procarbazine in patients with glioblastoma multiforme at first relapse. *Br J Cancer* 2000, 83: 588-93.
47. Brandes, A.A., Tosoni, A., Amista, P., Nicolardi, L., Grosso, D., Berti, F., et al., How effective is BCNU in recurrent glioblastoma in the modern era? A phase II trial. *Neurology* 2004, 63: 1281-4.
48. Scoccianti, S., Detti, B., Sardaro, A., Iannalfi, A., Meattini, I., Leonulli, B.G., et al., Second-line chemotherapy with fotemustine in temozolomide-pretreated patients with relapsing glioblastoma: a single institution experience. *Anticancer Drugs* 2008, 19: 613-20.
49. Brandes, A.A., Tosoni, A., Franceschi, E., Blatt, V., Santoro, A., Faedi, M., et al., Fotemustine as second-line treatment for recurrent or progressive glioblastoma after concomitant and/or adjuvant temozolomide: a phase II trial of Gruppo Italiano Cooperativo di Neuro-Oncologia (GICNO). *Cancer Chemother Pharmacol* 2009, 64: 769-75.
50. Addeo, R., Caraglia, M., De Santi, M.S., Montella, L., Abbruzzese, A., Parlato, C., et al., A new schedule of fotemustine in temozolomide-pretreated patients with relapsing glioblastoma. *J Neurooncol* 2011, 102: 417-24.
51. Fabrini, M.G., Silvano, G., Lolli, I., Perrone, F., Marsella, A., Scotti, V., et al., A multi-institutional phase II study on second-line Fotemustine chemotherapy in recurrent glioblastoma. *J Neurooncol* 2009, 92: 79-86.
52. Batchelor, T.T., Mulholland, P., Neyns, B., Nabors, L.B., Campone, M., Wick, A., et al., Phase III randomized trial comparing the efficacy of cediranib as monotherapy, and in combination with lomustine, versus lomustine alone in patients with recurrent glioblastoma. *J Clin Oncol* 2013, 31: 3212-8.
53. Wick, W., Puduvalli, V.K., Chamberlain, M.C., van den Bent, M.J., Carpentier, A.F., Cher, L.M., et al., Phase III study of enzastaurin compared with lomustine in the treatment of recurrent intracranial glioblastoma. *J Clin Oncol* 2010, 28: 1168-74.
54. Taal, W., Oosterkamp, H.M., Walenkamp, A.M., Dubbink, H.J., Beerepoot, L.V., Hanse, M.C., et al., Single-agent bevacizumab or lomustine versus a combination of bevacizumab plus lomustine in patients with recurrent glioblastoma (BELOB trial): a randomised controlled phase 2 trial. *Lancet Oncol* 2014, 15: 943-53.
55. Brandes, A.A., Carpentier, A.F., Kesari, S., Sepulveda, J.M., Wheeler, H., Chinot, O., et al., A phase II study of galunisertib monotherapy or galunisertib plus lomustine compared to lomustine monotherapy in recurrent glioblastoma *J Clin Oncol* 33 2015, 33: (suppl; abstr 2014).
56. Wick, W., Brandes, A., Gorlia, T., Bendszus, M., Sahm, F., Taal, W., et al., Phase III trial exploring the combination of bevacizumab and lomustine in patients with first recurrence of a glioblastoma: the EORTC 26101 trial. *Neuro Oncol* 2015, 17: (suppl 5; LB05)

57. van den Bent, M.J., Brandes, A.A., Rampling, R., Kouwenhoven, M.C., Kros, J.M., Carpentier, A.F., et al., Randomized phase II trial of erlotinib versus temozolomide or carmustine in recurrent glioblastoma: EORTC brain tumor group study 26034. *J Clin Oncol* 2009, 27: 1268-74.
58. Weathers, S., Han, S., Liu, D., Conrad, C.A., Gilbert, M.R., Loghin, M.E., et al., A randomized phase II trial of standard dose bevacizumab versus low dose bevacizumab plus lomustine (CCNU) in adults with recurrent glioblastoma. *J Clin Oncol* 2015, 33: (suppl; abstr 2005).
59. Soffietti, R., Trevisan, E., Bertero, L., Cassoni, P., Morra, I., Fabrini, M.G., et al., Bevacizumab and fotemustine for recurrent glioblastoma: a phase II study of AINO (Italian Association of Neuro-Oncology). *J Neurooncol* 2014, 116: 533-41.
60. Brandes, A.A., Tosoni, A., Basso, U., Reni, M., Valduga, F., Monfardini, S., et al., Second-line chemotherapy with irinotecan plus carmustine in glioblastoma recurrent or progressive after first-line temozolomide chemotherapy: a phase II study of the Gruppo Italiano Cooperativo di Neuro-Oncologia (GICNO). *J Clin Oncol* 2004, 22: 4779-4786.
61. Prados, M.D., Yung, W.K., Fine, H.A., Greenberg, H.S., Junck, L., Chang, S.M., et al., Phase 2 study of BCNU and temozolomide for recurrent glioblastoma multiforme: North American Brain Tumor Consortium study. *Neuro Oncol* 2004, 6: 33-7.
62. Gaviani, P., Salmaggi, A., and Silvani, A., Combined chemotherapy with temozolomide and fotemustine in recurrent glioblastoma patients. *J Neurooncol* 2011, 104: 617-8.
63. Esteller, M., Garcia-Foncillas, J., Andion, E., Goodman, S.N., Hidalgo, O.F., Vanaclocha, V., et al., Inactivation of the DNA-repair gene MGMT and the clinical response of gliomas to alkylating agents. *N Engl J Med* 2000, 343: 1350-4.
64. Nagane, M., Kobayashi, K., Ohnishi, A., Shimizu, S., and Shiokawa, Y., Prognostic significance of O6-methylguanine-DNA methyltransferase protein expression in patients with recurrent glioblastoma treated with temozolomide. *Jpn J Clin Oncol* 2007, 37: 897-906.
65. Brandes, A.A., Ermani, M., Basso, U., Amista, P., Berti, F., Scienza, R., et al., Temozolomide as a second-line systemic regimen in recurrent high-grade glioma: a phase II study. *Ann Oncol* 2001, 12: 255-7.
66. Brandes, A.A., Ermani, M., Basso, U., Paris, M.K., Lumachi, F., Berti, F., et al., Temozolomide in patients with glioblastoma at second relapse after first line nitrosourea-procarbazine failure: a phase II study. *Oncology* 2002, 63: 38-41.
67. Chan, D.T., Poon, W.S., Chan, Y.L., and Ng, H.K., Temozolomide in the treatment of recurrent malignant glioma in Chinese patients. *Hong Kong Med J* 2005, 11: 452-6.

68. Perry, J.R., Belanger, K., Mason, W.P., Fulton, D., Kavan, P., Easaw, J., et al., Phase II trial of continuous dose-intense temozolomide in recurrent malignant glioma: RESCUE study. *J Clin Oncol* 2010, 28: 2051-7.
69. Brada, M., Stenning, S., Gabe, R., Thompson, L.C., Levy, D., Rampling, R., et al., Temozolomide versus procarbazine, lomustine, and vincristine in recurrent high-grade glioma. *J Clin Oncol* 2010, 28: 4601-8.
70. Reardon, D.A., Nabors, L.B., Mason, W.P., Perry, J.R., Shapiro, W., Kavan, P., et al., Phase I/randomized phase II study of afatinib, an irreversible ErbB family blocker, with or without protracted temozolomide in adults with recurrent glioblastoma. *Neuro Oncol* 2015, 17: 430-439.
71. Weller, M., Tabatabai, G., Kastner, B., Felsberg, J., Steinbach, J.P., Wick, A., et al., MGMT Promoter Methylation Is a Strong Prognostic Biomarker for Benefit from Dose-Intensified Temozolomide Rechallenge in Progressive Glioblastoma: The DIRECTOR Trial. *Clin Cancer Res* 2015, 21: 2057-64.
72. Desjardins, A., Reardon, D.A., Coan, A., Marcello, J., Herndon, J.E., Bailey, L., et al., Bevacizumab and daily temozolomide for recurrent glioblastoma. *Cancer* 2012, 118: 1302-1312.
73. Verhoeff, J.J., Lavini, C., van Linde, M.E., Stalpers, L.J., Majoie, C.B., Reijneveld, J.C., et al., Bevacizumab and dose-intense temozolomide in recurrent high-grade glioma. *Ann Oncol* 2010, 21: 1723-7.
74. Reardon, D.A., Desjardins, A., Peters, K., Gururangan, S., Sampson, J., Rich, J.N., et al., Phase II study of metronomic chemotherapy with bevacizumab for recurrent glioblastoma after progression on bevacizumab therapy. *J Neurooncol* 2011, 103: 371-9.
75. Sepulveda, J.M., Belda-Iniesta, C., Gil-Gil, M., Perez-Segura, P., Berrocal, A., Reynes, G., et al., A phase II study of feasibility and toxicity of bevacizumab in combination with temozolomide in patients with recurrent glioblastoma. *Clin Transl Oncol* 2015.
76. Groves, M.D., Puduvalli, V.K., Gilbert, M.R., Levin, V.A., Conrad, C.A., Liu, V.H., et al., Two phase II trials of temozolomide with interferon-alpha2b (pegylated and non-pegylated) in patients with recurrent glioblastoma multiforme. *Br J Cancer* 2009, 101: 615-20.
77. Reardon, D.A., Vredenburgh, J.J., Desjardins, A., Peters, K., Gururangan, S., Sampson, J.H., et al., Effect of CYP3A-inducing anti-epileptics on sorafenib exposure: results of a phase II study of sorafenib plus daily temozolomide in adults with recurrent glioblastoma. *J Neurooncol* 2011, 101: 57-66.
78. Quinn, J.A., Jiang, S.X., Reardon, D.A., Desjardins, A., Vredenburgh, J.J., Rich, J.N., et al., Phase II trial of temozolomide plus o6-benzylguanine in adults with recurrent, temozolomide-resistant malignant glioma. *J Clin Oncol* 2009, 27: 1262-7.

79. Reardon, D.A., Quinn, J.A., Rich, J.N., Desjardins, A., Vredenburgh, J., Gururangan, S., et al., Phase I trial of irinotecan plus temozolomide in adults with recurrent malignant glioma. *Cancer* 2005, 104: 1478-86.
80. Silvani, A., Eoli, M., Salmaggi, A., Lamperti, E., Maccagnano, E., Broggi, G., et al., Phase II trial of cisplatin plus temozolomide, in recurrent and progressive malignant glioma patients. *J Neurooncol* 2004, 66: 203-8.
81. Brandes, A.A., Basso, U., Reni, M., Vastola, F., Tosoni, A., Cavallo, G., et al., First-line chemotherapy with cisplatin plus fractionated temozolomide in recurrent glioblastoma multiforme: a phase II study of the Gruppo Italiano Cooperativo di Neuro-Oncologia. *J Clin Oncol* 2004, 22: 1598-604.
82. Chua, S.L., Rosenthal, M.A., Wong, S.S., Ashley, D.M., Woods, A.M., Dowling, A., et al., Phase 2 study of temozolomide and Caelyx in patients with recurrent glioblastoma multiforme. *Neuro Oncol* 2004, 6: 38-43.
83. Gan, H., Fichtel, L., Lassman, A., Merrell, R., van den Bent, M., Kumthekar, P., et al., A phase 1 study evaluating ABT-414 in combination with temozolomide (TMZ) for subjects with recurrent or unresectable glioblastoma (GBM). *J Clin Oncol* 2015, 32:5s (suppl; abstr 2021)
84. Field, K.M., Simes, J., Nowak, A.K., Cher, L., Wheeler, H., Hovey, E.J., et al., Randomized phase 2 study of carboplatin and bevacizumab in recurrent glioblastoma. *Neuro Oncol* 2015, 17:1504-13
85. Friedman, H.S., Prados, M.D., Wen, P.Y., Mikkelsen, T., Schiff, D., Abrey, L.E., et al., Bevacizumab alone and in combination with irinotecan in recurrent glioblastoma. *J Clin Oncol* 2009, 27: 4733-40.
86. Kreisl, T.N., Kim, L., Moore, K., Duic, P., Royce, C., Stroud, I., et al., Phase II trial of single-agent bevacizumab followed by bevacizumab plus irinotecan at tumor progression in recurrent glioblastoma. *J Clin Oncol* 2009, 27: 740-5.
87. Raizer, J.J., Grimm, S., Chamberlain, M.C., Nicholas, M.K., Chandler, J.P., Muro, K., et al., A phase 2 trial of single-agent bevacizumab given in an every-3-week schedule for patients with recurrent high-grade gliomas. *Cancer* 2010, 116: 5297-305.
88. Nagane, M., Nishikawa, R., Narita, Y., Kobayashi, H., Takano, S., Shinoura, N., et al., Phase II study of single-agent bevacizumab in Japanese patients with recurrent malignant glioma. *Jpn J Clin Oncol* 2012, 42: 887-95.
89. Puduvalli, V., Wu, J., Yuan, Y., Armstrong, T., Groves, M.D., Raizer, J., et al., Brain Tumor Trials Collaborative Bayesian Adaptive Randomized Phase II trial of bevacizumab plus vorinostat versus bevacizumab alone in adults with recurrent glioblastoma (BTTC-1102). *J Clin Oncol* 2015, 33: (suppl; abstr 2012).
90. Galanis, E., Keith Anderson, S., Anastasiadis, P., Tran, D.D., Jeyapalan, S.A., Anderson, D.M., et al., NCCTG N0872 (Alliance): A randomized placebo-controlled phase II trial of bevacizumab plus dasatinib in patients with recurrent glioblastoma (GBM). *J Clin Oncol* 2015, 33: (suppl; abstr 2004).

91. Stark-Vance, V., Bevacizumab and CPT-11 in the treatment of relapsed malignant glioma. *Neuro Oncol* 2005, 7: 369 (abstr 342).
92. Vredenburgh, J.J., Desjardins, A., Herndon, J.E., 2nd, Marcello, J., Reardon, D.A., Quinn, J.A., et al., Bevacizumab plus irinotecan in recurrent glioblastoma multiforme. *J Clin Oncol* 2007, 25: 4722-9.
93. Vredenburgh, J.J., Desjardins, A., Herndon, J.E., 2nd, Dowell, J.M., Reardon, D.A., Quinn, J.A., et al., Phase II trial of bevacizumab and irinotecan in recurrent malignant glioma. *Clin Cancer Res* 2007, 13: 1253-9.
94. Gilbert, M.R., Wang, M., Aldape, K.D., Lassman, A.B., Sorensen, A.G., Mikkelsen, T., et al., RTOG 0625: A phase II study of bevacizumab with irinotecan in recurrent glioblastoma (GBM) *J Clin Oncol* 2009, 27: (suppl; abstr 2011).
95. Moller, S., Grunnet, K., Hansen, S., Schultz, H., Holmberg, M., Sorensen, M., et al., A phase II trial with bevacizumab and irinotecan for patients with primary brain tumors and progression after standard therapy. *Acta Oncol* 2012, 51: 797-804.
96. Hasselbalch, B., Lassen, U., Hansen, S., Holmberg, M., Sorensen, M., Kosteljanetz, M., et al., Cetuximab, bevacizumab, and irinotecan for patients with primary glioblastoma and progression after radiation therapy and temozolomide: a phase II trial. *Neuro Oncol* 2010, 12: 508-16.
97. Reardon, D.A., Desjardins, A., Peters, K.B., Gururangan, S., Sampson, J.H., McLendon, R.E., et al., Phase II study of carboplatin, irinotecan, and bevacizumab for bevacizumab naive, recurrent glioblastoma. *J Neurooncol* 2012, 107: 155-64.
98. Reardon, D.A., Desjardins, A., Vredenburgh, J.J., Gururangan, S., Sampson, J.H., Sathornsumetee, S., et al., Metronomic chemotherapy with daily, oral etoposide plus bevacizumab for recurrent malignant glioma: a phase II study. *Br J Cancer* 2009, 101: 1986-94.
99. Lassen, U., Sorensen, M., Gaziel, T.B., Hasselbalch, B., and Poulsen, H.S., Phase II study of bevacizumab and temsirolimus combination therapy for recurrent glioblastoma multiforme. *Anticancer Res* 2013, 33: 1657-60.
100. Sathornsumetee, S., Desjardins, A., Vredenburgh, J.J., McLendon, R.E., Marcello, J., Herndon, J.E., et al., Phase II trial of bevacizumab and erlotinib in patients with recurrent malignant glioma. *Neuro Oncol* 2010, 12: 1300-10.
101. Galanis, E., Anderson, S.K., Lafky, J.M., Uhm, J.H., Giannini, C., Kumar, S.K., et al., Phase II study of bevacizumab in combination with sorafenib in recurrent glioblastoma (N0776): a north central cancer treatment group trial. *Clin Cancer Res* 2013, 19: 4816-23.
102. Lee, E.Q., Reardon, D.A., Schiff, D., Drappatz, J., Muzikansky, A., Grimm, S.A., et al., Phase II study of panobinostat in combination with bevacizumab for recurrent glioblastoma and anaplastic glioma. *Neuro Oncol* 2015.

103. Ghiaieddin, A., Reardon, D., Massey, W., Mannerino, A., Lipp, E. S., Herndon, J.E. and al., Phase II study of bevacizumab and vorinostat for recurrent glioblastoma. *J Clin Oncol* 2015, 33: (suppl; abstr 2034).
104. Bennouna, J., Sastre, J., Arnold, D., Osterlund, P., Greil, R., Van Cutsem, E., et al., Continuation of bevacizumab after first progression in metastatic colorectal cancer (ML18147): a randomised phase 3 trial. *Lancet Oncol* 2013, 14: 29-37.
105. Hovey, E., Field, K., Rosenthal, M., Nowak, A., Cher, L., Wheeler, H., Continuing or ceasing bevacizumab at disease progression: Results from the CABARET study, a prospective randomized phase II trial in patients with recurrent glioblastoma. *J Clin Oncol* 2015, 33: (suppl; abstr 2003).
106. Batchelor, T.T., Duda, D.G., di Tomaso, E., Ancukiewicz, M., Plotkin, S.R., Gerstner, E., et al., Phase II study of cediranib, an oral pan-vascular endothelial growth factor receptor tyrosine kinase inhibitor, in patients with recurrent glioblastoma. *J Clin Oncol* 2010, 28: 2817-23.
107. Lassen, U., Chinot, O.L., McBain, C., Mau-Sorensen, M., Larsen, V.A., Barrie, M., et al., Phase 1 dose-escalation study of the antiplacental growth factor monoclonal antibody RO5323441 combined with bevacizumab in patients with recurrent glioblastoma. *Neuro Oncol* 2015. 17:1007-15.
108. Kreisl, T., Lassman, A., Mischel, P., Rosen, N., Scher, H., Teruya-Feldstein, J., et al., A pilot study of everolimus and gefitinib in the treatment of recurrent glioblastoma (GBM). *J Neurooncol* 2009, 92: 99-105.
109. Lassen, U., Mau-Soerensen, M., L. Kung, A., Wen, P.Y., Lee, E.Q., Plotkin, S.R., et al., A phase 2 study on efficacy, safety and intratumoral pharmacokinetics of oral selinexor (KPT-330) in patients with recurrent glioblastoma (GBM). *J Clin Oncol* 33, 2015 2015, 33.
110. Pitz, M.W., Eisenhauer, E.A., MacNeil, M.V., Thiessen, B., Easaw, J.C., Macdonald, D.R., et al., Phase II study of PX-866 in recurrent glioblastoma. *Neuro Oncol* 2015.
111. Lassman, A.B., Pugh, S.L., Gilbert, M.R., Aldape, K.D., Geinoz, S., Beumer, J.H., et al., Phase 2 trial of dasatinib in target-selected patients with recurrent glioblastoma (RTOG 0627). *Neuro Oncol* 2015.
112. Reardon, D.A., Fink, K.L., Mikkelsen, T., Cloughesy, T.F., O'Neill, A., Plotkin, S., et al., Randomized phase II study of cilengitide, an integrin-targeting arginine-glycine-aspartic acid peptide, in recurrent glioblastoma multiforme. *J Clin Oncol* 2008, 26: 5610-7.
113. Galanis, E., Jaeckle, K.A., Maurer, M.J., Reid, J.M., Ames, M.M., Hardwick, J.S., et al., Phase II trial of vorinostat in recurrent glioblastoma multiforme: a north central cancer treatment group study. *J Clin Oncol* 2009, 27: 2052-8.
114. Friday, B.B., Anderson, S.K., Buckner, J., Yu, C., Giannini, C., Geoffroy, F., et al., Phase II trial of vorinostat in combination with bortezomib in recurrent

glioblastoma: a north central cancer treatment group study. *Neuro Oncol* 2012, 14: 215-21.

115. Peereboom, D.M., Ahluwalia, M.S., Ye, X., Supko, J.G., Hilderbrand, S.L., Phuphanich, S., et al., NABTT 0502: a phase II and pharmacokinetic study of erlotinib and sorafenib for patients with progressive or recurrent glioblastoma multiforme. *Neuro Oncol* 2013, 15: 490-6.
116. Sampson, J., Vlahovic, G., Sahebjam, S., Padula Omuro, A.M., Baehring, J.M., Hafler, D.A., et al., Preliminary safety and activity of nivolumab and its combination with ipilimumab in recurrent glioblastoma (GBM): CHECKMATE-143. *J Clin Oncol* 2015, 33: (suppl; abstr 3010).
117. Hunn, M.K., Bauer, E., Wood, C.E., Gasser, O., Dzhelali, M., Ancelet, L.R., et al., Dendritic cell vaccination combined with temozolomide retreatment: results of a phase I trial in patients with recurrent glioblastoma multiforme. *J Neurooncol* 2015, 121: 319-29.
118. Schijns, V.E., Pretto, C., Devillers, L., Pierre, D., Hofman, F.M., Chen, T.C., et al., First clinical results of a personalized immunotherapeutic vaccine against recurrent, incompletely resected, treatment-resistant glioblastoma multiforme (GBM) tumors, based on combined allo- and auto-immune tumor reactivity. *Vaccine* 2015, 33: 2690-6.
119. Bloch, O., Crane, C.A., Fuks, Y., Kaur, R., Aghi, M.K., Berger, M.S., et al., Heat-shock protein peptide complex-96 vaccination for recurrent glioblastoma: a phase II, single-arm trial. *Neuro Oncol* 2014, 16: 274-9.
120. Reardon, D.A., Schuster, J., Tran, D.D., Fink, K., Nabors, L.B., Li, G., et al., ReACT: Overall survival from a randomized phase II study of rindopepimut (CDX-110) plus bevacizumab in relapsed glioblastoma. *J Clin Oncol* 2015, 33: (suppl; abstr 2009).
121. Stupp, R., Wong, E.T., Kanner, A.A., Steinberg, D., Engelhard, H., Heidecke, V., et al., NovoTTF-100A versus physician's choice chemotherapy in recurrent glioblastoma: a randomised phase III trial of a novel treatment modality. *Eur J Cancer* 2012, 48: 2192-202.
122. Rieger, J., Bahr, O., Maurer, G.D., Hattingen, E., Franz, K., Brucker, D., et al., ERGO: a pilot study of ketogenic diet in recurrent glioblastoma. *Int J Oncol* 2014, 44: 1843-52.
123. Khan, R.B., Raizer, J.J., Malkin, M.G., Bazylewicz, K.A., and Abrey, L.E., A phase II study of extended low-dose temozolomide in recurrent malignant gliomas. *Neuro Oncol* 2002, 4: 39-43.
124. Wick, W., Steinbach, J.P., Kuker, W.M., Dichgans, J., Bamberg, M., and Weller, M., One week on/one week off: a novel active regimen of temozolomide for recurrent glioblastoma. *Neurology* 2004, 62: 2113-5.
125. Brandes, A.A., Tosoni, A., Cavallo, G., Bertorelle, R., Gioia, V., Franceschi, E., et al., Temozolomide 3 weeks on and 1 week off as first-line therapy for

recurrent glioblastoma: phase II study from gruppo italiano cooperativo di neuro-oncologia (GICNO). *Br J Cancer* 2006, 95: 1155-60.

126. Kong, D.S., Lee, J.I., Kim, W.S., Son, M.J., Lim do, H., Kim, S.T., et al., A pilot study of metronomic temozolomide treatment in patients with recurrent temozolomide-refractory glioblastoma. *Oncol Rep* 2006, 16: 1117-21.
127. Wick, A., Felsberg, J., Steinbach, J.P., Herrlinger, U., Platten, M., Blaschke, B., et al., Efficacy and tolerability of temozolomide in an alternating weekly regimen in patients with recurrent glioma. *J Clin Oncol* 2007, 25: 3357-61.
128. Balmaceda, C., Peereboom, D., Pannullo, S., Cheung, Y.K., Fisher, P.G., Alavi, J., et al., Multi-institutional phase II study of temozolomide administered twice daily in the treatment of recurrent high-grade gliomas. *Cancer* 2008, 112: 1139-46.
129. Berrocal, A., Perez Segura, P., Gil, M., Balana, C., Garcia Lopez, J., Yaya, R., et al., Extended-schedule dose-dense temozolomide in refractory gliomas. *J Neurooncol* 2010, 96: 417-22.
130. Kong, D.-S., Lee, J.-I., Kim, J.H., Kim, S.T., Kim, W.S., Suh, Y.-L., et al., Phase II trial of low-dose continuous (metronomic) treatment of temozolomide for recurrent glioblastoma. *Neuro Oncol* 2010, 12: 289-296.
131. Santoni, M., Paccapelo, A., Burattini, L., Bianconi, M., Cardinali, M., Fabbietti, L., et al., Protracted low doses of temozolomide for the treatment of patients with recurrent glioblastoma: A phase II study. *Oncol Lett* 2012, 4: 799-801.
132. Norden, A.D., Lesser, G.J., Drappatz, J., Ligon, K.L., Hammond, S.N., Lee, E.Q., et al., Phase 2 study of dose-intense temozolomide in recurrent glioblastoma. *Neuro Oncol* 2013, 15: 930-935.
133. Han, S.J., Rolston, J.D., Molinaro, A.M., Clarke, J.L., Prados, M.D., Chang, S.M., et al., Phase II trial of 7 days on/7 days off temozolomide for recurrent high-grade glioma. *Neuro Oncol* 2014, 16: 1255-62.
134. Wen, P.Y., Prados, M., Schiff, D., Reardon, D.A., Cloughesy, T., Mikkelsen, T., et al., Phase II study of XL184 (BMS 907351), an inhibitor of MET, VEGFR2, and RET, in patients (pts) with progressive glioblastoma (GB). *J Clin Oncol* 2010, 28: (suppl; abstr 2006).
135. Wen, P.Y., American Society of Clinical Oncology 2010: report of selected studies from the CNS tumors section. *Expert Rev Anticancer Ther* 2010, 10: 1367-9.
136. Iwamoto, F.M., Lamborn, K.R., Robins, H.I., Mehta, M.P., Chang, S.M., Butowski, N.A., et al., Phase II trial of pazopanib (GW786034), an oral multi-targeted angiogenesis inhibitor, for adults with recurrent glioblastoma (North American Brain Tumor Consortium Study 06-02). *Neuro Oncol* 2010, 12: 855-861.
137. Stupp, R., Tosoni, A., Bromberg, J.E., Hau, P., Campone, M., Gijtenbeek, J., et al., Sagopilone (ZK-EPO, ZK 219477) for recurrent glioblastoma. A phase II

multicenter trial by the European Organisation for Research and Treatment of Cancer (EORTC) Brain Tumor Group. *Ann Oncol* 2011, 22: 2144-9.

138. de Groot, J.F., Lamborn, K.R., Chang, S.M., Gilbert, M.R., Cloughesy, T.F., Aldape, K., et al., Phase II study of aflibercept in recurrent malignant glioma: a North American Brain Tumor Consortium study. *J Clin Oncol* 2011, 29: 2689-95.
139. Wen, P.Y., Schiff, D., Cloughesy, T.F., Raizer, J.J., Laterra, J., Smitt, M., et al., A phase II study evaluating the efficacy and safety of AMG 102 (rilotumumab) in patients with recurrent glioblastoma. *Neuro Oncol* 2011, 13: 437-46.
140. Kreisl, T.N., McNeill, K.A., Sul, J., Iwamoto, F.M., Shih, J., and Fine, H.A., A phase I/II trial of vandetanib for patients with recurrent malignant glioma. *Neuro Oncol* 2012, 14: 1519-1526.
141. Pan, E., Yu, D., Yue, B., Potthast, L., Chowdhary, S., Smith, P., et al., A prospective phase II single-institution trial of sunitinib for recurrent malignant glioma. *J Neurooncol* 2012, 110: 111-8.
142. Muhic, A., Poulsen, H., Sorensen, M., Grunnet, K., and Lassen, U., Phase II open-label study of nintedanib in patients with recurrent glioblastoma multiforme. *J Neurooncol* 2013, 111: 205-212.
143. Reardon, D.A., Groves, M.D., Wen, P.Y., Nabors, L., Mikkelsen, T., Rosenfeld, S., et al., A Phase I/II Trial of Pazopanib in Combination with Lapatinib in Adult Patients with Relapsed Malignant Glioma. *Clin Cancer Res* 2013, 19: 900-908.
144. Kreisl, T., Smith, P., Sul, J., Salgado, C., Iwamoto, F., Shih, J., et al., Continuous daily sunitinib for recurrent glioblastoma. *J Neurooncol* 2013, 111: 41-48.
145. Chamberlain, M.C., Grimm, S., Phuphanich, S., Recht, L., Zhu, J.Z., Kim, L., et al., A phase 2 trial of verubulin for recurrent glioblastoma: a prospective study by the Brain Tumor Investigational Consortium (BTIC). *J Neurooncol* 2014, 118: 335-43.
146. Gilbert, M.R., Kuhn, J., Lamborn, K.R., Lieberman, F., Wen, P.Y., Mehta, M., et al., Cilengitide in patients with recurrent glioblastoma: the results of NABTC 03-02, a phase II trial with measures of treatment delivery. *J Neurooncol* 2012, 106: 147-53.

Figure 1: Approach for individualized treatment decisions in patients with glioblastoma.

Continuous arrows indicate evidence-based current clinical practice. Dashed arrows represent possibilities of individual decision-making which has still to be confirmed.

Abbreviations: CCNU, lomustine; KPS, Karnofsky performance scale; RT, radiotherapy; TMZ, temozolomide; TMZ/RT→TMZ, radiotherapy with concomitant and maintenance TMZ

